

Sex-specific Effects of the DAF-12 Steroid Receptor on Aging in *Caenorhabditis elegans*

DIANA McCULLOCH AND DAVID GEMS

Centre for Research on Ageing, Department of Biology, University College London, London WC1E 6BT, United Kingdom

ABSTRACT: Sex differences in longevity and aging are seen throughout the animal kingdom. These are likely to result, in part, from sex differences in endocrinology. In the nematode *Caenorhabditis elegans*, males are the longer-lived sex. Here we explore the possibility that sex differences in insulin/insulin-like growth factor 1 (IGF-1) and steroid endocrinology contribute to this sex difference in aging by studying *C. elegans* populations in liquid culture. We report that in hermaphrodite populations, mutational loss of the DAF-12 steroid receptor affected life span as in previous plate-culture studies: mutant longevity is suppressed in a weak *daf-2* insulin/IGF-1 receptor mutant but enhanced in a stronger *daf-2* mutant. However, in males, mutation of *daf-12* had little effect on aging in either weak or strong *daf-2* mutants. Moreover, while mutation of *daf-12* marginally reduced life span in *daf-2*(+) hermaphrodites, as in plate-cultured populations, it did not in *daf-2*(+) males. These results could imply that in *C. elegans*, as in mammals, sex differences in steroid endocrinology contribute to sex differences in aging.

KEYWORDS: *Caenorhabditis elegans*; aging; gender; steroid; insulin/IGF-1 signaling

INTRODUCTION

A salient feature of human aging is that women age slightly more slowly than men. In developed country this leads to a gender gap in mean life span of some 4–6 years.¹ Sex differences in aging are typical of animal species, but their biological basis remains poorly understood.² Many aspects of sexual dimorphism are specified by sex differences in endocrine function, and this may apply to aging too. For example, the shorter life span of men is thought to arise, in part, from the effects of testicular hormones (e.g., the steroid hormone testosterone) on behavior and on cardiovascular health.¹ While in many animal

Address for correspondence: David Gems, Department of Biology, University College London, London WC1E 6BT, UK. Voice: (44) (0) 20 7679 4381; fax: (44) (0) 20 7679 7096. david.gems@ucl.ac.uk

Ann. N.Y. Acad. Sci. 1119: 253–259 (2007). © 2007 New York Academy of Sciences.
doi: 10.1196/annals.1404.018

species, females are longer lived, in the nematode *Caenorhabditis elegans* the male is the longer-lived sex, living some 20% longer than hermaphrodites.³

Several endocrine pathways are known to influence longevity and aging in *C. elegans*. Insulin/insulin-like growth factor 1 (IGF-1) signaling shortens life span,⁴ and mutation of the *daf-2* insulin/IGF-1 receptor gene can more than double adult life span.⁵ Life span is also modulated by bile acid-like steroid hormones which act via the nuclear receptor DAF-12.^{6,7} The interplay between insulin-like and steroid pathways in the control of aging is complex. In adult hermaphrodites, mutational inactivation of *daf-12* slightly shortens life span in a *daf-2(+)* background, partially suppresses the longevity phenotype of weaker *daf-2* mutants, but enhances that of more severe *daf-2* mutants.^{8,9} Thus, the DAF-12-mediated steroid pathway can increase or decrease life span, depending on the genetic context.

In this study, we explore the possibility that steroid hormone biology in *C. elegans* might, like that of higher animals, be sexually dimorphic and contribute to sex differences in aging. To do this, we have directly compared the effect on life span in males and hermaphrodites of mutation of *daf-12*, either alone or in combination with mutations affecting *daf-2*. We report that the effects of *daf-12* on aging differ substantially between the sexes.

EXPERIMENTAL METHODS

Nematodes were raised on NGM agar seeded with *Escherichia coli* OP50 bacteria, as previously described.¹⁰ Measurement of life span in *C. elegans* males is complicated by male–male interactions, which shorten life span, and frequent escape of males from agar plates.³ Therefore, for this study, life span was measured in monoxenic liquid culture with *E. coli* OP50, with animals maintained individually in microtiter wells, as previously described.¹¹

The three mutant alleles studied were as follows: *daf-12(m20)* results in a dauer defective phenotype and has a stop codon near the 5' end of the open reading frame, and therefore approximates to a null allele⁶; *daf-2(m41)* and *daf-2(e1370)* are both hypomorphic alleles resulting in constitutive formation of dauer larvae and increased adult longevity—*m41* is a less pleiotropic (class 1) allele with a missense mutation in the Cys-rich region of the receptor, while *e1370* is a more pleiotropic (class 2) allele with a missense mutation in the kinase region of the receptor.^{12,13}

RESULTS AND DISCUSSION

We compared the effects on aging in both sex of mutations of *daf-12*, alone or in combination with mutation of *daf-2*. In hermaphrodites, *daf-12(m20)* alone caused a marginal nonsignificant reduction in life span (TABLE 1, FIG. 1).

TABLE 1. Effect of *daf-12* on life span in both sexes and in *daf-2(+)* and *daf-2* mutant backgrounds

Genotype/ gender	Median life span (days) ± 95% C.I.	% Effect of <i>daf-12</i> on median	Maximum life span (days)	% Effect of <i>daf-12</i> on maximum	N*	P†
+ H ^a	17.7 (19.0, 16.5)	—	23.0	—	165 (200)	—
+ M ^b	19.8 (21.3, 18.3)	—	29.0	—	135 (211)	—
<i>daf-12(m20)</i> H	16.0 (17.5, 15.0)	-10	20.0	-13	88 (151)	0.56
<i>daf-12(m20)</i> M	23.5 (24.5, 22.1)	+19	29.5	+2	131 (152)	0.07, 0.0001
<i>daf-2(m41)</i> H	31.8 (36.0, 28.8)	—	51.5	—	44 (80)	—
<i>daf-2(m41)</i> M	29.0 (32.0, 26.5)	—	41.5	—	59 (80)	—
<i>daf-2(m41); daf-12</i> H †	18.0 (20.0, 18.0)	-43	22.0	-57	37 (80)	<0.0001
<i>daf-2(m41); daf-12</i> M	30.0 (33.5, 28.0)	+3	44.0	+6	87 (90)	0.017, 0.74
<i>daf-2(e1370)</i> H	29.3 (32.0, 27.3)	—	45.0	—	124 (332)	—
<i>daf-2(e1370)</i> M	53.0 (56.0, 50.8)	—	105.0	—	209 (262)	—
<i>daf-2(e1370); daf-12</i> H †	54.0 (75.0, 43.5)	+84	113.5	+152	30 (72)	<0.0001
<i>daf-2(e1370); daf-12</i> M †	64.0 (83.0, 55.0)	+21	124.0	+18	40 (72)	0.18

^aHermaphrodite; ^bMale; * Senescent deaths (starting population); †Probability that survival curves of a strain with and without *daf-12(m20)* differ by random chance (log rank test). Multiple significance values represent results from trials performed at separate times. ‡Results from only one replicate because of bacterial contamination in two other replicates. Trials performed at 22.5 °C.




FIGURE 1. Survival curves showing effect of *daf-12(m20)* on class 1 *daf-2(m41)* and class 2 *daf-2(e1370)* hermaphrodite and male survival (22.5 °C). N2 (wild type) and *daf-12(m20)* are shown for comparison. P = probability that survival of *daf-2(rf)* and *daf-2(rf)*; *daf-12(m20)* differ by random chance (log rank test).

		Phenotype promoted by DAF-12	
		Hermaphrodites	Males
+		Longevity	Aging
	<i>daf-2(m41)</i>	Longevity	(Aging)
	<i>daf-2(e1370)</i>	Aging	No effect

FIGURE 2. Summary of the inferred effects of the DAF-12 steroid receptor on longevity and aging. In all three genetic contexts, the effects of mutation of *daf-12* on life span differ between males and hermaphrodites.

This is consistent with earlier studies using agar plates where *daf-12* significantly shortened hermaphrodite life span.^{8,9} *daf-12* also suppressed the increased longevity (Age phenotype) of the less pleiotropic (class 1) mutant *daf-2(m41)* but enhanced that of the more pleiotropic (class 2) mutant *daf-2(e1370)* (TABLE 1, FIG. 1). These results also tally with earlier studies of plate-cultured populations, where longevity of class 1 *daf-2* alleles was suppressed and that of class 2 *daf-2* alleles enhanced.^{8,9} The only difference from earlier studies is that here *daf-12* fully suppressed *daf-2(m41)* Age, whereas previously only partial suppression was seen. However, in a subsequent test of the effect of *daf-12* on class 1 *daf-2* life span in liquid culture, we saw only partial suppression of Age (data not shown). Thus, the use of liquid culture rather than plate culture does not appear to markedly alter the effects on aging of mutation of *daf-12* or *daf-2*, alone or in combination.

In males, *daf-12* did not shorten life span but instead produced a slight increase that was significant in one trial, and almost significant in another (FIG. 1, TABLE 1). In other trials, we also saw a significant increase in male life span resulting from the null allele *daf-12(rh61rh411)* (data not shown). Thus, the DAF-12 nuclear receptor weakly promotes longevity in hermaphrodites but weakly antagonizes it in males.

Both *daf-2(m41)* and *daf-2(e1370)* increased life span in males in liquid culture (FIG. 1, TABLE 1). The sex difference in longevity was suppressed by *daf-2(m41)*, something not seen in studies of another class 1 mutant, *daf-2(m577)*

on plates,³ By contrast, longevity was greatly increased in *daf-2(e1370)* males, as previously seen in plate-cultured populations of this genotype.³ Unexpectedly, addition of *daf-12* had only minor effects on life span in *daf-2* males (FIG. 1, TABLE 1). *daf-12* marginally increased *daf-2* male life span, but the effects were very small compared to the effects seen in hermaphrodites. These slight increases reached statistical significance in one of two trials with *daf-2(m41)*, but were not significant for *daf-2(e1370)*.

In summary, our results show that the effects of *daf-12* on aging differ between the sexes in all three backgrounds tested: *daf-2(+)*, *daf-2(m41)*, and *daf-2(e1370)* (findings summarized in FIG. 2). In a wild-type background, the DAF-12 receptor weakly promotes longevity in the hermaphrodite but aging in the male. In a weak (class 1) *daf-2* mutant background, DAF-12 promotes longevity in hermaphrodites but marginally promotes aging in males. In a stronger (class 2) *daf-2* mutant background, DAF-12 promotes longevity in hermaphrodites but has little effect in males.

These findings demonstrate sexual dimorphism in the effects on aging of the DAF-12 nuclear receptor. This could be determined in a number of ways, including sex differences in levels or tissue localization of DAF-12 protein levels, or of cofactors that interact with DAF-12, or in production or structure of the steroid ligand that controls DAF-12 activity. Whether nematodes are, like mammals, sexually dimorphic in terms of the steroid hormones that they secrete, remains to be explored.

ACKNOWLEDGMENTS

We thank Scott Pletcher for assistance with data analysis. This work was supported by the Biotechnology and Biological Science Research Council (UK) and the Wellcome Trust.

REFERENCES

1. SMITH, D.W.E. 1993. Women live longer than men. *In* Human Longevity. Pp. 78–98. Oxford University Press. New York, Oxford.
2. MAY, R. 2007. Gender, immunity and the regulation of longevity. *BioEssays*. **29**: 795–802.
3. GEMS, D. & D.L. RIDDLE. 2000. Genetic, behavioral and environmental determinants of male longevity in *Caenorhabditis elegans*. *Genetics* **154**: 1597–1610.
4. KENYON, C. 2005. The plasticity of aging: insights from long-lived mutants. *Cell* **120**: 449–460.
5. KENYON, C. *et al.* 1993. A *C. elegans* mutant that lives twice as long as wild type. *Nature* **366**: 461–464.
6. ANTEBI, A. *et al.* 2000. *daf-12* encodes a nuclear receptor that regulates the dauer diapause and developmental age in *C. elegans*. *Genes Dev.* **14**: 1512–1527.

7. GERISCH, B. *et al.* 2007. A bile acid-like steroid modulates *Caenorhabditis elegans* lifespan through nuclear receptor signaling. *Proc. Natl. Acad. Sci. USA* **104**: 5014–5019.
8. GEMS, D. *et al.* 1998. Two pleiotropic classes of *daf-2* mutation affect larval arrest, adult behavior, reproduction and longevity in *Caenorhabditis elegans*. *Genetics* **150**: 129–155.
9. LARSEN, P.L., P.S. ALBERT & D.L. RIDDLE. 1995. Genes that regulate both development and longevity in *Caenorhabditis elegans*. *Genetics* **139**: 1567–1583.
10. SULSTON, J. & J. HODGKIN. 1988. Methods. *In* The nematode *Caenorhabditis elegans*. W.B. Wood, Ed.: 587–606. Cold Spring Harbor, NY.
11. MCCULLOCH, D. & D. GEMS. 2003. Evolution of male longevity advantage in nematodes. *Aging Cell* **2**: 165–173.
12. KIMURA, K.D. *et al.* 1997. *daf-2*, an insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science* **277**: 942–946.
13. YU, H. & P. LARSEN. 2001. DAF-16-dependent and independent expression targets of DAF-2 insulin receptor-like pathway in *Caenorhabditis elegans* include FKBP. *J. Mol. Biol.* **314**: 1017–1028.