

## Support for people bereaved by suicide

This section focuses on supporting people bereaved by suicide with the aim of reducing the possible negative effects of a suicide. This activity is commonly referred to as 'postvention'.

### Aims of postvention

An ability to draw on knowledge of the aims of postvention, which include, for example:
reducing the incidence of suicide and improving mental health in those who have been bereaved
reducing the stigma and isolation experienced by those who have been bereaved by suicide
offering bespoke support for those bereaved by suicide (such as support through the legal process)
signposting to appropriate sources of support
reducing the risk of contagion or the emergence of suicide clusters

### Knowledge of bereavement

An ability to draw on knowledge that the experience of bereavement will be influenced by many factors, including cultural norms, belief systems, faith systems and life experiences

An ability to draw on knowledge:

of psychological models that describe the process of grieving as a series of phases, the order and timing of which will vary from person to person, but which broadly include:

initial shock and denial

acceptance of the reality of the loss

experiencing the pain and distress associated with the loss

adjusting to a new reality in which the person who has died is no longer present

that it is usual for people to oscillate between a focus on loss and some degree of avoidance

that rather than grieving leading to a 'detachment' from the person who has died, it usually involves adjusting and redefining the relationship to them, allowing for a continued sense of connection and an enduring bond

## Knowledge about the nature of bereavement after a death by suicide

An ability to draw on knowledge that bereavement after a suicide is often different to bereavement after a death from other causes, arising from survivors' perceptions of, and preoccupation with, issues such as	
	the degree to which the suicide was seen as an understandable choice (e.g. suicide in the context of a terminal illness as contrasted to a death that is completely unexpected and not readily explicable)
	the extent to which the death is perceived as preventable, and the survivors' or other people's role and responsibility in relation to it (associated with feelings such as guilt, blame or anger)
	a sense of stigma or shame (which may lead to a withdrawal from potential sources of support)
	a feeling of isolation stemming from a loss that they feel others may not understand
An ability to draw on knowledge that survivors may experience trauma, for example, arising from:	
	ruminating about the intensity of the difficulties that the person who has died may have been experiencing
	direct exposure, such as witnessing the suicide itself (e.g. discovering the body), visiting the scene of death, being exposed to artefacts related to the suicide, or hearing detailed accounts of the death (e.g. in a coroner's court)
	imagined exposure (creating in the imagination, and ruminating on, a mental image of the way in which the person died and the suffering this entailed)
An ability to draw on knowledge that whilst grieving, close relatives of the person who has died will face unfamiliar legal and practical issues (such as the police investigation, identification of a body, the coronial process, delayed registration of death and the person's affairs) resulting from the nature of the death and its unexpectedness	
An ability to draw on knowledge of common reactions in people bereaved after a death by suicide, such as	
	cognitive and emotional reactions
	shock and disbelief
	anguish, longing and searching
	shame, and self-blame
	despair, depression and sadness
	a sense of abandonment and rejection by the person
	blaming others and assigning responsibility for the death
	trying to make sense of the death in order to restore a sense of order
	guilt (about actions taken or not taken and failure to save the person)
	anger, often related to feelings of guilt, blame, abandonment and a sense of preventability
	relief, often accompanied by confusion and guilt (e.g. if the suicide comes after a prolonged period of conflict or need for support)
	searching for answers (e.g. a preoccupation with learning about what led to the death, difficulty living with ambiguity surrounding the death)
	fear and anxiety (e.g. that another person close to them may take their own life)
	physical reactions (often associated with anxiety and/or depression) such as:
	stomach pains, tightness in the chest, sensitivity to noise, breathlessness, disturbed sleep, poor concentration, lack of energy
	symptoms of trauma, such as recurring nightmares or flashbacks

## Supporting people after a death by suicide

### Psychological support for people who have been bereaved by suicide

An ability to help to validate and to 'normalise' the person's experiences and to 'normalise' their reactions (e.g. through listening and through psychoeducation about bereavement)
An ability to allow the person to detail the circumstances of the death and its aftermath
An ability to help the person discuss their emotional reactions to the death and thereby:
come to terms with the reality of the death and the meanings associated with it
acknowledge, accept and find ways to manage their emotions
adjust to the inevitable changes associated with the suicide
rebuild connections to the person who has died that are based on reality (rather than idealised or excessively denigratory)
An ability to identify whether the bereaved person is themselves at risk of suicide and if so to implement a safety plan
An ability to identify resources and sources of support with legal and practical issues
An ability to enable access to support groups and psychological services for people bereaved by suicide

### Peer support groups

An ability for peer facilitators to:
draw on their own lived experience of bereavement by suicide
listen to the stories of others without being immobilised by their own grief
acknowledge that people will have different ways of coping
work with group members whose cultural background and experience is different from their own
respond appropriately to group members with a different suicide experience from their own
work collaboratively with a co-facilitator
maintain the confidentiality of the group
manage challenging group situations in an appropriate manner (e.g. making sure that certain individuals do not dominate the group)
access support if they are finding it difficult to cope with the cumulative effect of supporting others in distress

## Organisational competences relevant to postvention

An ability to draw on knowledge about the structure and content of postvention services set out in public health guidance and the recommendations contained in national suicide prevention strategies
An ability to gather and interpret local data on death by suicide that can be used to inform the need for postvention, for example by:
collating demographic data on suicide deaths
identifying patterns or trends in data on suicide deaths
identifying geographical areas of concern or other evidence of clusters
An ability to identify current local provision of postvention and the availability of funding for services
An ability to identify and contract with organisations/agencies with the capacity and ability to set-up and to deliver postvention services
An ability to identify the required characteristic of postvention services, for example:
identifying the recipients of services, such as:
adults only, adults and children, children and young people only
those with close connections to the person who has died, such as next of kin and/or close family members
non-family who were close to the person, such as friends
people who knew (but who had variable levels of contact with) the person who has died, such as school friends or work colleagues
identifying the ways in which the characteristics of recipients of services will influence service specifications
identifying what the service offers to recipients, for example:
information and signposting only
outreach to people who are bereaved
practical support (e.g. with funerals, legal processes)
individual clinical support
peer support
group support facilitated by professionals or by peers
referral pathways to other services
annual remembrance services
An ability to audit the effectiveness of the postvention service (e.g. through feedback from service users and those offering postvention services, or a requirement for formal reporting on uptake and use)

## Support for people within an organisation after a suicide

This section focuses on the organisational competences required to coordinate a response to a suicide that impacts on an organisation (such as a school or college, healthcare organisation or a workplace).

It identifies the competences associated with offering psychological support to affected individuals or the workplace as a whole, intended to reduce the possible negative effects of a suicide. This activity is commonly referred to as 'postvention'

### Knowledge of postvention

An ability to draw on knowledge that postvention refers to a range of interventions intended to ameliorate the negative impact of a death by suicide

An ability to draw on knowledge that when a suicide impacts an organisation (such as a workplace or school), postvention can be offered both to individuals and at an organisational level

An ability to draw on knowledge that the stance taken by the organisation's leaders sets the tone for the ways in which people in the organisation respond

### Instituting postvention

An ability for those in a position of leadership to convey empathy for those who have been bereaved, and to recognise:

the potential impact on others and indicate the importance of self-care

that the responses of members of the organisation will differ depending, for example, on their relationship to, and prior involvement with, the person who has died, and on their own history and experience of bereavement

that the responses of members of the organisation will differ depending on their own personal history and particularly their own experience of bereavement

that psychological reactions to the death will vary from person to person, both immediately and over time (e.g. shock, guilt, blaming others, anger, sadness)

that distress can be expressed in different ways, both explicitly and implicitly (through withdrawal, absenteeism or presenteeism)

that the level of support offered both to individuals and across the organisation will need to reflect an appraisal of the difficulties that need to be addressed

An ability for those in a position of leadership to recognise the impact of the death on themselves and to recognise their own need for support

An ability to draw on knowledge of basic principles for intervening after a suicide, namely:

that all deaths should be approached in the same way regardless of cause (in order to avoid stigmatising a death by suicide)

that because of the risk of suicide 'contagion', care should be taken not to glamorise or romanticise the person who has died or provide excessive detail about the method of suicide (such as the means or precise location of the death)

that the organisation should broadly indicate the likely reasons for suicide (e.g. that the person had psychological difficulties that may or may not have been apparent to others) and highlight that the causes of suicide are complex

that help should be made available for people who have similar feelings or are in distress

## Communicating information about the death

An ability to coordinate organisational communications about the death (e.g. being clear about who is responsible and the medium through which information is relayed)

An ability to draw on knowledge that it is helpful for communications about the death to be clear about the cause of death, for example:

making a full disclosure where the family or carers have consented for it to be known that the person died by suicide

where the family or carers are unwilling for this to be disclosed, managing this appropriately (e.g. indicating that the family or carers do not wish the cause of death to be known, but also facilitating discussion if there are rumours about suicide)

## Interventions

An ability to offer organisation-based interventions that can help:

to normalise reactions to the death by discussing the grieving process and the ways in which this might manifest over time, and the fact that there is no 'right' way of grieving

create opportunities to share stories, thoughts and memories

encourage discussion of the feelings engendered by the death (particularly expressions of guilt, anger or abandonment)

convey the need for self-care

An ability to make practical arrangements for workplace interventions, for example:

establishing debriefing meetings for groups and/or individuals, led by appropriately trained facilitators

varying working practices so as to allow time for meetings to take place

ensuring that there are follow-up meetings in place

An ability to ensure that there is long-term provision of support so as to recognise that feelings about the death can re-emerge (e.g. around the anniversary of the death, or around significant events that signal the absence of the person who has died)

An ability to draw on knowledge that most people will adapt to the death without professional support (reflecting their 'natural' resilience), but that some may experience more profound emotional reactions (such as complicated grief, trauma or suicidal ideation)

an ability for the organisation to have systems in place that identify people who are having significant difficulty coping and which can direct them towards appropriate sources of support

An ability to consider whether it is appropriate for the organisation to conduct a memorial event (with the aim of helping people to acknowledge the death and share their grief)

an ability to ensure that any ceremony strikes the right balance between commemorating but not memorialising (and potentially glamorising) the person who has died

## Judging when to end postvention

An ability to balance the need for changes in usual patterns of activity to accommodate reactions to the death against the need to re-establish the usual functioning of the organisation, and to:

judge the pace of a return in relation to an assessment of the degree to which the organisation and individuals within it have had time to adjust to the death

accommodate the needs of people who are not ready to return to their usual pattern of functioning