

Algorithm 1: Assessment

Using an ABCDE (Airway, Breathing, Circulation, Disability, Exposure) approach, assess whether the patient is hypovolaemic and needs fluid resuscitation
 Assess volume status taking into account clinical examination, trends and context. Indicators that a patient may need fluid resuscitation include: systolic BP <100mmHg; heart rate >90bpm; capillary refill >2s or peripheries cold to touch; respiratory rate >20 breaths per min; NEWS ≥5; 45° passive leg raising suggests fluid responsiveness.

Yes

Algorithm 2: Fluid Resuscitation

Initiate treatment

- Identify cause of deficit and respond.
- Give a fluid bolus of 500 ml of crystalloid (containing sodium in the range of 130–154 mmol/l) over 15 minutes.

Reassess the patient using the ABCDE approach
Does the patient still need fluid resuscitation? Seek expert help if unsure

Yes No

Does the patient have signs of shock?

Yes No

>2000 ml given?

Yes No

Give a further fluid bolus of 250–500 ml of crystalloid

Assess the patient's likely fluid and electrolyte needs

- History: previous limited intake, thirst, abnormal losses, comorbidities.
- Clinical examination: pulse, BP, capillary refill, JVP, oedema (peripheral/pulmonary), postural hypotension.
- Clinical monitoring: NEWS, fluid balance charts, weight.
- Laboratory assessments: FBC, urea, creatinine and electrolytes.

Can the patient meet their fluid and/or electrolyte needs orally or enterally?

Yes

Ensure nutrition and fluid needs are met
 Also see [Nutrition support in adults](#) (NICE clinical guideline 32).

No

Does the patient have complex fluid or electrolyte replacement or abnormal distribution issues?
 Look for existing deficits or excesses, ongoing abnormal losses, abnormal distribution or other complex issues.

Yes

Algorithm 4: Replacement and Redistribution

Existing fluid or electrolyte deficits or excesses
 Check for:

- dehydration
- fluid overload
- hyperkalaemia/hypokalaemia

Estimate deficits or excesses.

Ongoing abnormal fluid or electrolyte losses
 Check ongoing losses and estimate amounts. Check for:

- vomiting and NG tube loss
- biliary drainage loss
- high/low volume ileal stoma loss
- diarrhoea/excess colostomy loss
- ongoing blood loss, e.g. melaena
- sweating/fever/dehydration
- pancreatic/jejunal fistula/stoma loss
- urinary loss, e.g. post AKI polyuria.

Redistribution and other complex issues
 Check for:

- gross oedema
- severe sepsis
- hypernatraemia/hyponatraemia
- renal, liver and/or cardiac impairment.
- post-operative fluid retention and redistribution
- malnourished and refeeding issues

Seek expert help if necessary and estimate requirements.

Prescribe by adding to or subtracting from routine maintenance, adjusting for all other sources of fluid and electrolytes (oral, enteral and drug prescriptions)

Monitor and reassess fluid and biochemical status by clinical and laboratory monitoring


Algorithm 3: Routine Maintenance

Give maintenance IV fluids
 Normal daily fluid and electrolyte requirements:

- 25–30 ml/kg/d water
- 1 mmol/kg/day sodium, potassium, chloride
- 50–100 g/day glucose (e.g. glucose 5% contains 5 g/100ml).

Reassess and monitor the patient
 Stop IV fluids when no longer needed. Nasogastric fluids or enteral feeding are preferable when maintenance needs are more than 3 days.

Algorithm 1: Assessment



Using an ABCDE (Airway, Breathing, Circulation, Disability, Exposure) approach, assess whether the patient is hypovolaemic and needs fluid resuscitation

Assess volume status taking into account clinical examination, trends and context. Indicators that a patient may need fluid resuscitation include: systolic BP <100mmHg; heart rate >90bpm; capillary refill >2s or peripheries cold to touch; respiratory rate >20 breaths per min; NEWS \geq 5; 45° passive leg raising suggests fluid responsiveness.

Yes

Algorithm 2: Fluid Resuscitation

Initiate treatment

- Identify cause of deficit and respond.
- Give a fluid bolus of 500 ml of crystalloid (containing sodium in the range of 130–154 mmol/l) over 15 minutes.

Reassess the patient using the ABCDE approach
Does the patient still need fluid resuscitation? Seek expert help if unsure

Yes

No

Does the patient have signs of shock?

Yes

No

Assess the patient's likely fluid and electrolyte needs (Refer algorithm 1 box 3)


>2000 ml given?


Yes

Seek expert help

No

Give a further fluid bolus of 250–500 ml of crystalloid





Algorithm 4: Replacement and Redistribution

