

Peripheral Inflammation Acutely Impairs Human Spatial Memory via Actions on Medial Temporal Lobe Glucose Metabolism

Neil A. Harrison, Christian F. Doeller, Valerie Voon, Neil Burgess, and Hugo D. Critchley

Background: Inflammation impairs cognitive performance and is implicated in the progression of neurodegenerative disorders. Rodent studies demonstrated key roles for inflammatory mediators in many processes critical to memory, including long-term potentiation, synaptic plasticity, and neurogenesis. They also demonstrated functional impairment of medial temporal lobe (MTL) structures by systemic inflammation. However, human data to support this position are limited.

Methods: Sequential fluorodeoxyglucose positron emission tomography together with experimentally induced inflammation was used to investigate effects of a systemic inflammatory challenge on human MTL function. Fluorodeoxyglucose positron emission tomography scanning was performed in 20 healthy participants before and after typhoid vaccination and saline control injection. After each scanning session, participants performed a virtual reality spatial memory task analogous to the Morris water maze and a mirror-tracing procedural memory control task.

Results: Fluorodeoxyglucose positron emission tomography data demonstrated an acute reduction in human MTL glucose metabolism after inflammation. The inflammatory challenge also selectively compromised human spatial, but not procedural, memory; this effect that was independent of actions on motivation or psychomotor response. Effects of inflammation on parahippocampal and rhinal glucose metabolism directly mediated actions of inflammation on spatial memory.

Conclusions: These data demonstrate acute sensitivity of human MTL to mild peripheral inflammation, giving rise to associated functional impairment in the form of reduced spatial memory performance. Our findings suggest a mechanism for the observed epidemiologic link between inflammation and risk of age-related cognitive decline and progression of neurodegenerative disorders including Alzheimer's disease.

Key Words: Alzheimer's disease, imaging, inflammation, memory, parahippocampus, PET

Although previously considered an immune-privileged site, it is now clear that the immune system plays an integral role in many fundamental neuronal processes, including long-term potentiation (LTP) (1,2), synaptic plasticity (3), and neurogenesis (4), that are critical to learning and memory. In health, immune mechanisms regulate each of these processes and assist in the remodeling of neural circuits that promote learning and memory (5). However, during systemic infection or injury (6), this positive regulatory function is disrupted, resulting in acute memory impairments: When inflammation is severe, cognitive impairment may become persistent (7), and when chronic inflammation is present, age-related cognitive impairment is accelerated (8). Inflammation may drive the rapid progression of neurodegenerative diseases such as Alzheimer's disease (9).

From the Department of Psychiatry (NAH, HDC), Brighton and Sussex Medical School, and Sackler Centre for Consciousness Science (NAH, HDC), University of Sussex, Falmer, United Kingdom; Donders Institute for Brain, Cognition and Behaviour (CFD), Radboud University Nijmegen, The Netherlands; Department of Psychiatry (VV), University of Cambridge, Cambridge, United Kingdom; and Institutes of Cognitive Neuroscience and Neurology (NB), University College London, London, United Kingdom.

Address correspondence to Neil A. Harrison, M.B.B.S., Ph.D., Clinical Imaging Sciences Centre, Brighton and Sussex Medical School, University of Sussex, Falmer, BN1 9RR, United Kingdom; E-mail: n.harrison@bsms.ac.uk.

Received Oct 16, 2013; revised and accepted Jan 7, 2014.

0006-3223/\$36.00

<http://dx.doi.org/10.1016/j.biopsych.2014.01.005>

Structures in the medial temporal lobe (MTL) appear to be particularly sensitive to effects of inflammation. This increased sensitivity may be related to their relatively high receptor and messenger RNA expression for proinflammatory cytokines (10,11) and their neural connectivity to regions such as the insula (12) that support cortical representations of peripheral inflammatory states (13). Rodent studies emphasized the role of the hippocampus; direct administration of inflammatory cytokines, particularly interleukin (IL)-1, into the hippocampus selectively impaired spatial and contextual memory processes, including radial arm and Morris water maze performance, and contextual, but not auditory-cued, fear conditioning (5,14,15). Similarly, overexpression of IL-1 messenger RNA within the hippocampus is associated with delayed acquisition of spatial memory on the Morris water maze task (14). For synaptic plasticity underlying the encoding and recall of memories, LTP is arguably the key neuronal mechanism. The cytokine IL-1 compromises both hippocampal and dentate gyrus LTP (1,17,18) and may mediate both age-dependent decreases in LTP (19) and the modulation of LTP by A β amyloid (20). Cytokine-induced inhibition of neurogenesis within the dentate gyrus is also alleviated by the microglial inhibitor minocycline (4). Together, these data highlight the central action of inflammatory mediators (cytokines such as IL-1) on MTL-dependent memory processes.

Inflammatory challenges administered outside the central nervous system also induce IL-1 expression within brain regions, including the MTL (21). Peripherally induced inflammation also replicates many of the direct actions of inflammatory cytokines on MTL-dependent memory (5,22,23). There are numerous mechanisms through which peripheral inflammation can engender changes in cytokine levels within sensitive brain regions. Circulating cytokines may be actively transported across the blood-brain barrier

(24) or activate microglia via the circumventricular organs (25) and vascular endothelium (26). However, local synthesis of IL-1 is suggested by the rapid upregulation of IL-1 α and IL-1 β gene expression and the central predominance of the short half-life IL-1 isoform in the context of mild systemic inflammatory challenge (21). Vagus nerve afferents show sensitivity to peripheral cytokines (27) and mild inflammatory challenge (28) indicating an additional neurally mediated immune-brain pathway. Central vagus nerve targets show enhanced activity within 2–3 hours of peripheral inflammatory challenge in both rodents and humans (29,30). Electrical stimulation of vagus nerve afferents results in a rapid increase in IL-1 β expression within the hippocampus (31). Humoral and neurally mediated routes may communicate peripheral inflammatory responses centrally to regions supporting memory processes.

These data from animal studies suggest mechanisms to account for human epidemiologic data linking increased peripheral inflammation to accelerated cognitive aging and neurodegeneration. However, it is unknown whether systemic inflammation modulates MTL function in humans. We used an experimental inflammatory model, typhoid vaccination, together with sequential fluorodeoxyglucose (FDG) positron emission tomography (PET) scanning to quantify hypothesized effects of peripheral inflammation on human MTL function and spatial memory. In 20 healthy participants, three FDG-PET scans were performed immediately before and 4 hours and 8 hours after typhoid vaccination or control (saline) injection (Figure 1). After each of the first two scanning sessions, participants performed a spatial memory task in which they learned and then recalled the identity and location of two sets of 16 objects positioned within a virtual reality environment. This virtual reality task is analogous to the Morris water maze (32), which is sensitive to inflammatory effects on object-location accuracy in rodents, and to the hidden tracer task, which is sensitive to lesions in discrete MTL structures in humans (33). Recall of the spatial location and identity of both sets of objects was tested again after the third scan to investigate differential effects of inflammation on early encoding and later consolidation processes. Participants also performed a mirror-tracing procedural memory task to test general effects of inflammation on psychomotor responses and motor learning.

Methods and Materials

Participants

We recruited 20 healthy male nonsmokers (mean age, 24.7 \pm 6.8 years old) and screened them for relevant physical or

psychiatric illness; all were medication-free. Volunteers who had received typhoid vaccine within 3 years or other vaccine within 6 months were excluded. Participants were advised to avoid caffeinated beverages, alcohol, high-fat meals, and excessive exercise for 24 hours and steroidal or nonsteroidal drugs for 2 weeks before testing. All participants fasted for 8 hours and consumed only water until study completion. Written informed consent was obtained from all participants, and procedures were approved by the Brighton East National Research Ethics Committee.

Study Design

A randomized, double-blind, repeated measures crossover design was used in which all participants underwent three FDG-PET imaging sessions each separated by 4 hours. After each of the first two scanning sessions, participants randomly received intramuscular injections of either 0.025 mg *Salmonella typhi* vaccine (Typhim Vi; Aventis Pasteur MSD Ltd., Lyon, France) or 0.5 mL normal saline. Of participants, 13 were randomly assigned to the early inflammation group and received vaccination after the first PET scan 1, and 7 were randomly assigned to the late inflammation group and received vaccination after the second PET scan. This study design enabled us to control for nonspecific time effects as well as have sufficient participants ($n = 13$) scanned 8 hours after typhoid vaccination to test late effects of inflammation. After each scan, participants performed a laptop-based spatial memory task and a mirror-tracing procedural memory task that took 35 min to complete. Vaccination or saline injection was given after the PET scan immediately before memory testing; this was done to minimize an already long testing day. We are aware of no data to suggest that peripherally induced inflammation can impair memory at such a short latency, and if this were the case, it would increase the risk of false-negative rather than false-positive findings. A high-resolution inversion recovery echo planar image was obtained to aid image registration.

Inflammatory Model

We used a *S. typhi* vaccination model known to induce low-grade inflammation without body temperature change (34). Blood (10 mL) was drawn into ethylenediamine tetraacetic acid BD Vacutainer tubes (Franklin Lakes, New Jersey) and centrifuged at 1250 $\times g$ for 10 min, and plasma was removed, aliquoted, and frozen at -80°C . Plasma IL-6, IL-1 receptor antagonist, and tumor necrosis factor alpha were assessed using high-sensitivity enzyme-linked immunosorbent assays (R&D Systems, Abingdon, United Kingdom). Limits of detection were .039 pg/mL, 6.26 pg/mL,




Figure 1. Study timeline. All participants completed three fluorodeoxyglucose (FDG) positron emission tomography (PET) scans. Each scan was preceded by a blood draw and followed by a memory testing session. The “early” inflammation group received the typhoid vaccination after the first PET scan (and saline injection after the second PET scan), and the “late” inflammation group received the typhoid vaccination after the second PET scan (saline injection after the first scan). In the first two sessions (T1 and T2), participants encoded and then recalled the identity and spatial location of two sets of objects (object set 1 and set 2). In the third session (T3), participants recalled the identity and spatial location of the two sets of objects encoded at T1 and T2. The mirror-tracing task was performed at each of the three testing sessions. MRI, magnetic resonance imaging. (Photo credit, copyright WGBH Educational Foundation.)

and .038 pg/mL with associated intra-assay and interassay coefficients of variation of 7.4% and 7.8%, 5.3% and 8.6%, and 5.3% and 8.4%.

Spatial Memory Task

UnrealEngine2 Runtime software (Epic Games, Cary, North Carolina) was used to present a first-person perspective of a plane surrounded by a circular cliff (virtual diameter 180 m). Background mountains, clouds, and the sun (created using Terragen; Planetside Software, Cheshire, United Kingdom) projected at infinity were used to provide orientation cues. Two separate counterbalanced arenas with grassy or rocky planes and differently rendered mountains and clouds were used for the two encoding sessions. Participants explored the arena for 2–3 min using right-handed button presses to move forward, left, and right. Then 16 unique objects were sequentially presented within the arena, and participants were instructed to remember their identity and spatial location before picking them up. After all objects were acquired, participants performed free recall of the object identities. They then returned to the virtual reality environment. A picture of one object was presented, following which participants moved to where they thought the cued object had been presented, indicated by a button press, and recorded their confidence (range, 1–5) for this location. For the first two sessions (T1 and T2), the object was shown again in its correct position, and participants collected it by running over it. The next object was cued after a variable intertrial interval. On the third session (T3), object recall and relocation phases were completed for objects learned at T1 (set 1) and T2 (set 2). Performance was indexed by accuracy of object spatial location: mean (1/distance from true object location in virtual meters) and number of objects recalled.

Procedural Memory Task

Participants were asked to trace between two concentric five-pointed stars viewed in a mirror as quickly and accurately as possible. Both their hand and the concentric stars were obscured from direct view. Time taken to complete two trials was used as an index of performance.

Image Acquisition and Analysis

The PET scans (mean 155.3 ± 11.8 MBq FDG) were acquired for 35 min on a Siemens Biograph-64 PET-CT scanner (Siemens Healthcare, Erlangen, Germany) in three-dimensional dynamic acquisition mode. Participants lay supine with eyes open. Before each PET acquisition, a low-dose computed tomography scan (120 kVp, 10 mA) was acquired for attenuation correction. After correction for scatter, random effects, and effects of attenuation, images were reconstructed in 1-min windows using Siemens proprietary iterative three-dimensional reconstruction schema (21 iterations and 8 subsets). Individual 1-min scans were realigned and summed to produce a single 35-min activation scan per session, which was coregistered to subjects' structural magnetic resonance imaging scans and then spatially smoothed with an 8-mm full width at half maximum Gaussian kernel using standard SPM8 methods (Wellcome Trust Centre for Neuroimaging, Institute of Neurology, United College London, London, United Kingdom; <http://www.fil.ion.ucl.ac.uk/spm>).

Normalized images were included within a second-level flexible-factorial analysis of variance (ANOVA) (repeated factor time, baseline, 4 hours, 8 hours; between-subject factor group, early, late inflammation). Main effects of time and group and group \times time interaction were included in the model.

Normalization to a grand mean scaled value of 50 mL/100 g/min was applied, and global effects were included as nuisance covariates in the general linear model (analysis of covariance). Correlations between changes in resting glucose metabolism and object-location accuracy together with interactions with inflammatory status (modeled using a dummy variable) were investigated in a separate regression analysis.

Anatomic localization of MTL structures was based on Insausti *et al.* (35), Monte Carlo simulation (1000 iterations) was used to determine cluster extent thresholds for whole-brain correction at $p < .01$ (36), and a cluster threshold of >19 voxels was adopted. Regression analyses followed by Goodman test for mediation were used to investigate relationships between inflammatory challenge and changes in object-location accuracy and right parahippocampal activity Montreal Neurological Institute (24, –32, –32) between encoding sessions T1 and T2. Mediation analyses were performed using the interactive calculation tool for mediation tests (<http://quantpsy.org/sobel/sobel.htm>).

Results

Inflammatory Responses to Typhoid Vaccination

Cytokine analyses confirmed significantly higher circulating inflammatory cytokines at encoding session 2 (T2) compared with encoding session 1 (T1) in the early, but not late, inflammation group (Figure 2A): group \times time interaction IL-6 [$F_{1,18} = 6.91, p = .017$], IL-1 receptor antagonist [$F_{1,18} = 11.77, p = .003$]. A similar increase in IL-6 (2.82 pg/mL) was also observed after typhoid vaccination (between T2 and T3) in the late inflammation group (Figure S1 in Supplement 1). Both groups showed inflammation at T3 and absence of inflammation at baseline (T1) with only the early inflammation group inflamed at encoding session two (T2).

Effects of Acute Inflammation on Memory Performance

Immediate recall of object identity and location at T2 and T1 demonstrated a significant group \times encoding session interaction for object location but not object identity [$F_{1,17} = 5.01, p = .039$ and $F_{1,17} = .66, p = .43$]. Post hoc *t* tests also demonstrated a significantly greater reduction in proximity score across the two encoding sessions (T1 and T2) in the early inflammation group, who demonstrated inflammation at T2, compared with the late inflammation group, who did not demonstrate inflammation at this time point ($-.100$ vs. $.234$ m^{-1} ; $t_{17} = 2.24, p = .039$) (Figure 2B,C). Although performance increased from session T1 to session T2 in the late inflammation group given placebo at T1 (a practice effect), it decreased in the early inflammation group given vaccine, suggesting that inflammation impaired object-location encoding during the T2 session. This effect was maintained at the later recall session (T3) when both sets of objects were recalled and both groups demonstrated inflammation: group \times encoding session interaction [$F_{1,17} = 8.40, p = .01$]. Post hoc *t* test again demonstrated a significantly greater reduction in proximity score at T3 for objects encoded at T2 compared with objects encoded at T1 in the early compared with late inflammation group ($-.061$ vs. $.287$ m^{-1} ; $t_{17} = 2.90, p = .01$) (Figure 2E,F). The impairing effect of inflammation on encoding location of objects seen at T2 was preserved when these objects were later recalled at T3.

Performance on the mirror-tracing task revealed no significant group \times time interaction [$F_{1,18} = 1.00, p = .33$], although improved performance across time was observed in both groups




Figure 2. Cytokine and memory performance in the early and late inflamed groups. **(A)** Plasma interleukin-6 levels demonstrating a significantly greater increase in interleukin-6 (from session T1 to T2) in the early compared with late inflammation group. **(B)** Object location accuracy (proximity score) in units of 1/virtual meters at each encoding session (T1 and T2). Location accuracy decreased 4 hours after typhoid vaccination (early inflammation group) but increased 4 hours after placebo (late inflammation group). **(C)** Number of objects (maximum 16) correctly recalled during the two encoding sessions (T1 and T2). **(D)** Mean time taken to complete mirror tracing of a five-pointed star demonstrating a significant improvement across time in both groups. **(E)** Object location accuracy (proximity score) at the final session (T3) for objects encoded at T1 (set 1) and T2 (set 2). **(F)** Number of objects (maximum 16) correctly recalled at the final session (T3) for objects encoded at T1 (set 1) and T2 (set 2). Asterisk indicates statistical significance at $p < .05$. IL-6, interleukin-6.

[main effect of time $F_{2,18} = 23.58$, $p < .001$] (Figure 2D). These results suggest a selective action of inflammation on object-location memory that is not mediated via nonspecific effects on task motivation or response time.

To explore whether effects on object-location memory were mediated by actions at encoding or consolidation, we next performed a three-way ANOVA: group (early inflammation, late inflammation), encoding session (T1, T2), and recall session (T3 recall of first set, T3 recall of second set) with the prediction that actions at consolidation would be reflected by greater effects at late (T3) than early (T1 and T2) recall, owing to impaired consolidation in the early compared with the late inflammation group. This ANOVA confirmed the previously observed encoding session \times group interaction [$F_{1,17} = 4.44$, $p = .028$]. However, no additional recall session \times group or encoding session \times recall session \times group interactions were observed [$F_{1,17} = .84$, $p = .45$ and $F_{1,17} = .80$, $p = .47$] suggesting a predominant effect at encoding.

To address this situation further, we regressed immediate (T1 and T2) against late (T3) performance for both object sets with inclusion of a dummy variable encoding group membership. This regression demonstrated an anticipated strong dependence of late on early performance for both sets of objects [$F_{1,15} = 6.08$, $p = .026$ for object set 1 and $F_{1,15} = 29.67$, $p < .0001$ for object set 2] but no interaction with group [$F_{1,15} = 1.53$, $p = .26$ and $F_{1,15} = .77$, $p = .39$]. Finally, we performed a 2 (group) \times 2 (recall

session) ANOVA on performance at T3 corrected for T1 and T2 performance. This ANOVA failed to show a significant recall session \times group interaction [$F_{1,17} = 2.16$, $p = .16$]. Together, these analyses suggest a significant action of inflammation on early encoding and consolidation processes with little evidence to support additional effects on late consolidation processes. They also provide empiric support for a direct influence of inflammation on early encoding and consolidation mechanisms rather than nonspecific effects on motivation in which a greater decrement in performance at T3 compared with T2 would be expected in the late inflammation group (subjects demonstrated inflammation only at the later time point) compared with the early inflammation group (subjects demonstrated inflammation at both time points).

Effects of Acute Inflammation on Resting Brain Glucose Metabolism

Analysis of PET data (T1 and T2) in the early compared with the late inflammation group) demonstrated a reduction in glucose metabolism within a discrete cluster of regions focused on the right parahippocampal and perirhinal cortex 4 hours after inflammation compared with placebo (Table 1; Figure 3A). These regions all survived whole-brain correction at $p < .01$. This finding was replicated 4 hours after inflammatory challenge in the late compared with early inflammation group (T3 compared with T2) (Table 1; Figure 3B), robustly demonstrating acute sensitivity of MTL structures to peripheral inflammation.

Table 1. Brain Regions Showing an Acute Reduction in Resting Glucose Metabolism After Inflammatory Challenge

| Side | Region | Coordinates | Z Score | Cluster | $p_{\text{uncorrected}}$ | $p_{\text{corrected}}$ |
|--|----------------------------|---------------|---------|---------|--------------------------|------------------------|
| Inflammation-Induced Reductions in Glucose Metabolism (Early Inflammation Group) | | | | | | |
| R | Parahippocampus/perirhinal | (36 -28 -24) | 3.73 | 345 | <.001 | <.01 |
| R | Fusiform gyrus | (32 -54 -14) | 3.89 | 190 | <.001 | <.01 |
| R | Inferior temporal gyrus | (57 -18 -33) | 3.56 | 76 | <.001 | <.01 |
| R | Temporal pole | (57 11 -11) | 3.42 | 28 | <.001 | <.01 |
| Inflammation-Induced Reductions in Glucose Metabolism (Late Inflammation Group) | | | | | | |
| R | Entorhinal/perirhinal | (21 -19 -27) | 3.50 | 117 | <.001 | <.01 |
| L | Entorhinal/perirhinal | (-24 -13 -26) | 3.67 | 71 | <.001 | <.01 |
| R | Parahippocampus/perirhinal | (33 -24 -24) | 3.22 | 10 | <.001 | <.01 |

L, left; R, right.

To investigate whether this change in glucose metabolism between encoding sessions predicted changes in object-location accuracy, we next performed a regression analysis on the PET data (i.e., T1 and T2 metabolism vs. T1 and T2 accuracy) (Table 2). This analysis revealed striking correlations between activity change in bilateral parahippocampal and rhinal cortex and change in object-location accuracy across all participants (Figure 3C)—that is, there was a general relationship between change in parahippocampal and rhinal glucose metabolism and change in object-location accuracy. However, repetition of this analysis after inclusion of an interaction term coding group membership (early or late) also revealed a discrete contiguous

region within the right parahippocampal gyrus that mediated the detrimental effects of inflammation on object-location encoding (Figures 3D and 4). In other words, inflammation disrupted the relationship between parahippocampal metabolism and subsequent accuracy for object-location encoding. This interpretation was supported further by mediation analysis, which showed that inflammation induced changes in right parahippocampal glucose metabolism (T1 and T2) Montreal Neurological Institute (24, -21, -32) that significantly mediated effects of inflammation on object-location memory (T1 and T2 accuracy) (Goodman test = 3.58 [SE .74], $p < .00035$) (Figure 5).




Figure 3. Brain regions sensitive to acute inflammation and effects on object-location encoding. (A) Regions showing a greater reduction in glucose metabolism after inflammation compared with placebo between sessions 1 and 2. Contrast shown in T1 - T2 early group minus T1 - T2 late group. (B) Regions showing a greater reduction in glucose metabolism after inflammatory challenge in the late compared with early inflamed group between sessions 2 and 3. Contrast shown in T2 and T3 early group compared with T2 and T3 late group. The y axis in (A) and (B) shows estimated glucose metabolism in mL/100 g/min. (C) Regions showing a positive correlation between change in object-location accuracy and change in glucose metabolism between the two encoding sessions (T1 and T2) across all participants. (D) Medial temporal lobe region showing a significant group \times location accuracy interaction between the two encoding sessions (T1 and T2). The y axis shows change in glucose metabolism between T1 and T2 in mL/100 g/min. E, early inflammation group (received vaccine after first scan); L, late inflammation group (received vaccine after second scan 2).

Table 2. Brain Regions in Which Changes in Blood Glucose Metabolism Between the Two Encoding Sessions (T1 and T2) Predicted Associated Changes in Memory for Object Location

| Side | Region | Coordinates | Z Score | Cluster | $p_{\text{uncorrected}}$ | $p_{\text{corrected}}^a$ |
|---|----------------------------|---------------|---------|---------|--------------------------|--------------------------|
| Regions Showing Positive Correlations Across All Participants | | | | | | |
| R | Parahippocampus/perirhinal | (27 -22 -27) | 4.19 | 751 | <.001 | <.01 |
| L | Parahippocampus/perirhinal | (-29 -28 -26) | 3.38 | 80 | <.001 | <.01 |
| R | Precuneus | (15 -60 45) | 3.68 | 370 | <.001 | <.01 |
| L | Inferior parietal lobule | (-20 -51 54) | 3.65 | 829 | <.001 | <.01 |
| R | Inferior parietal lobule | (33 -46 56) | 3.31 | 99 | <.001 | <.01 |
| L | Supplementary motor area | (-15 -4 63) | 3.55 | 95 | <.001 | <.01 |
| L | Paracentral lobule | (-3 -31 63) | 3.36 | 221 | <.001 | <.01 |
| R | Mid-frontal gyrus | (35 3 46) | 3.21 | 26 | <.001 | <.01 |
| R | Mid-orbitofrontal gyrus | (48 50 -9) | 3.17 | 25 | <.001 | <.01 |
| Medial Temporal Lobe Region Showing Interaction with Group | | | | | | |
| R | Parahippocampus/perirhinal | (24 -21 -32) | 2.44 | 342 | <.05 | <.01 ^b |

L, left; R, right.

^a $p_{\text{corrected}}$ = cluster survives whole-brain correction.

^bSurvives correction for a medial temporal lobe region of interest.

Discussion

Systemic inflammation is associated with selective impairment in human spatial memory but not MTL-independent procedural memory (37). Deficits in spatial memory were observed for objects learned and recalled during systemic inflammation but not objects learned in the absence of inflammation and recalled under inflammatory conditions. This study suggests a predominant effect of inflammation on early encoding and consolidation

processes rather than late consolidation and demonstrates a relative absence of state-dependent effects. In rodent contextual fear conditioning paradigms (5,23), inflammation impairs spatial memory despite being induced after visuospatial information has been attended to (and encoded) suggesting that our data are likely mediated via an action on early consolidation processes. Although our analyses failed to demonstrate significant late consolidation effects, Figure 2B and E demonstrates a non-significant reduction in performance at T3 compared with T1 in the early (early T3 [set 1] compared with early T1) but not late inflammation group (late T3 [set 1] compared with late T1) consistent with a potential effect on late consolidation.

Resting glucose metabolism, particularly change in bilateral parahippocampus and perirhinal cortex metabolism immediately before task performance, predicted change in accuracy across encoding sessions across all participants (Figures 2C and 3). However, this relationship was critically modulated by systemic inflammation. Within 4 hours of inflammatory challenge, glucose metabolism decreased within perirhinal and entorhinal cortex and parahippocampus (Table 1). This effect was replicated in participants challenged after the first (Figure 3A) and second (Figure 3B) scanning sessions. A discrete subregion centered on the right parahippocampus also predicted and mediated inflammatory effects on subsequent object-location memory (Figures 3D and 5). Together, these data demonstrate sensitivity of human MTL structures, notably parahippocampus, to systemic




Figure 4. Right medial temporal lobe regions sensitive to inflammation, change in object-location accuracy, and interactions with inflammation. Cyan indicates regions showing a reduction in glucose metabolism after inflammation (T1 - T2 early group minus T1 - T2 late group). Yellow indicates regions showing a positive correlation between change in object-location accuracy (T1 and T2) and change in glucose metabolism (T1 and T2) across all participants. Red indicates area showing group × accuracy interaction—that is, the region mediating the effects of group membership (inflammation status) on change in accuracy for encoding object location.




Figure 5. Mediation analysis showing that the changes in parahippocampal glucose metabolism mediate the effects of inflammation on memory for object location. Path coefficients (standard error of path coefficients) are shown for each path of the mediation model.

inflammation and provide mechanistic insight relevant to a broader literature linking severe or chronic inflammation to the attrition of human memory.

Studies investigating effects of inflammation on rodent spatial memory to date have predominantly focused on actions on the hippocampus (1,5,15–17). We did not identify a major change in hippocampal glucose metabolism after inflammation or any association between hippocampal glucose metabolism and subsequent memory performance. Although null results are hard to interpret, learning object locations relative to the boundary in this task correlates with functional magnetic resonance imaging signal from both right hippocampal and parahippocampal regions (32), suggesting greater sensitivity to detect metabolic changes in the parahippocampus. In addition, there is good evidence to suggest strong parahippocampal involvement in this type of task. Neuropsychological studies show that human performance on homologues of the Morris water maze and direct tests of object-location memory can be more strongly dependent on right parahippocampal than hippocampal integrity (37–39). Studies demonstrate a central role for the right parahippocampus in human object-location memory and support our current finding of a critical role for the parahippocampus in mediating inflammation-induced spatial memory impairments.

Right parahippocampal activity during object-location encoding has also been shown to predict subsequent retrieval success with a spatial cue (40). In monkeys, one-trial memory for object-place associations (similar to our current task) appears to be critically dependent not on hippocampus but on posterior parahippocampus (41). The contribution of the parahippocampus to within-scene object location and context memory is also dissociable from the role of perirhinal cortex in object perception and memory (42–44). In rodents, perirhinal neurons respond selectively to objects and their previous occurrence (45,46) with selective lesions impairing performance on tasks requiring whole-object information (47). In contrast, rodents with postrhinal (parahippocampus) cortex lesions show impairment on tasks sensitive to object location but not identity (48). Similar functional distinctions between perirhinal and parahippocampal activity are also apparent in humans, with parahippocampal cortex active during object-location encoding and perirhinal cortex active to objects alone (49). Our data suggest that systemic inflammation may serve as a transient parahippocampal lesion resulting in a discrete impairment in object-location memory.

The cellular mechanisms mediating this selective impairment of human MTL function are unclear, although these mechanisms may be usefully informed by rodent studies. For example, IL-1 has been shown to reduce basal synaptic activity and synaptic transmission in a manner dependent on gamma-aminobutyric acid (50). It impairs LTP both dependent on and independent of *N*-methyl-D-aspartate (1,2) and can decrease LTP-associated glutamate release within the dentate gyrus (51). Although these effects are currently demonstrated only in rodent hippocampus, operation of either mechanism within human parahippocampus or perirhinal cortex could conceivably contribute to the observed reduction in glucose metabolism. Local or peripheral inflammation can also impair hippocampal neurogenesis in proportion to the associated increase in microglial activation (4). However, given the time course of this effect, it is unlikely to have contributed to our results. Perhaps more pertinent is the role of neurally mediated mechanisms. Peripheral inflammation has been shown to increase rapidly activity within vagus nerve projection areas in both rodents and humans (29,30), including insular cortex, a region that in primates has direct neural connectivity

to perirhinal and parahippocampal cortex (12), areas that provide the vast bulk of inputs into entorhinal cortex and the hippocampal formation. Electrical stimulation of vagus nerve afferents results in a rapid increase in IL-1 β expression within the hippocampus (31). Activation of neurally mediated immune-brain communicatory pathways may potentially modulate memory processes even in the absence of significant signaling of inflammation across the blood-brain barrier at the endothelium (26) or circumventricular organs (25).

A concern from rodent studies is that apparent effects of inflammation on learning and memory may be confounded by actions on psychomotor speed (52). We also previously reported psychomotor slowing after typhoid vaccination (53). However, our current data strongly argue against a purely psychomotor explanation for our effects. In particular, vaccination did not change time taken to relocate objects; the late inflammation group showed no decrement in recall performance at time three, and mirror-tracing task performance was unimpaired by inflammation. As such, our data support and reinforce the interpretation of rodent studies.

One unresolved question is why we did not observe an effect on object identity memory, especially given reduced glucose metabolism across an expansive MTL region encompassing perirhinal cortex. Although participants did not perform at ceiling, the relatively small number of exemplars may have reduced variability associated with this measure, and consequently it may have been insensitive to subtle changes in object identity memory. This interpretation is also suggested by data from studies that show more global reductions in memory after potent inflammatory challenges with lipopolysaccharide, which have evoked decreased immediate verbal recall of story items, immediate and delayed spatial figural features, and word list learning (54). In another study, using low-dose lipopolysaccharide challenge, declarative memory impairment was also inversely correlated with IL-6 levels (55).

Our study identifies a mechanism through which peripheral inflammation affects human spatial memory. This study has important implications for understanding how chronic inflammation exacerbates age-related cognitive decline and plausibly the increased risk of neurodegenerative disorders such as Alzheimer's disease. Increased inflammatory markers are observed in the MTL of patients with age-related cognitive decline and Alzheimer's disease (56). The profile of memory impairment observed in Alzheimer's disease—selective impairment of MTL-dependent memory including impaired spatial memory (57) with often striking preservation of procedural memory (58)—is similar to what we describe here. Nevertheless, it is uncertain whether they are the cause of cognitive symptoms or a consequence of a primary disease process. Increased circulating proinflammatory cytokines have been associated within an increased risk of cognitive decline in both cross-sectional and prospective epidemiologic studies (8). Similarly, acute infections requiring admission to the intensive care unit convey a significantly greater risk of subsequent cognitive decline compared with other causes of intensive care unit admission (7). In healthy middle-aged adults, levels of circulating inflammatory cytokines are linked to the volume of MTL structures, specifically hippocampus (59).

In conclusion, our data suggest that MTL structures are acutely sensitive to peripheral inflammation with consequent functional impairment. Peripheral inflammation results in an acute reduction in resting MTL glucose function associated with an acute decline in human spatial memory. This knowledge is motivation for

further investigation into the cognitive consequences of chronic or severe infections and inflammation.

This research was supported by Wellcome Trust Intermediate Clinical Fellowships awarded to NAH and VV, European Research Council support to CFD and HDC, Netherlands Organisation for Scientific Research support to CFD, and Medical Research Council (United Kingdom) and Wellcome Trust support to NB. We thank K. Miles for technical assistance with positron emission tomography imaging and E. Cooper and P. Ghezzi for assistance with enzyme-linked immunosorbent assays.

The authors report no biomedical financial interests or potential conflicts of interest.

Supplementary material cited in this article is available online at <http://dx.doi.org/10.1016/j.biopsych.2014.01.005>.

- Katsuki H, Nakai S, Hirai Y, Akaji K, Kiso Y, Satoh M (1990): Interleukin-1 β inhibits long-term potentiation in the CA3 region of mouse hippocampal slices. *Eur J Pharmacol* 181:323–332.
- Schneider H, Pitossi F, Balschun D, Wagner A, del Rey A, Besedovsky HO (1998): A neuromodulatory role of interleukin-1beta in the hippocampus. *Proc Natl Acad Sci U S A* 95:7778–7783.
- Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, *et al.* (2012): Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron* 74:691–705.
- Ekdahl CT, Claassen JH, Bonde S, Kokaia Z, Lindvall O (2003): Inflammation is detrimental for neurogenesis in adult brain. *Proc Natl Acad Sci U S A* 100:13632–13637.
- Yirmiya R, Goshen I (2011): Immune modulation of learning, memory, neural plasticity and neurogenesis. *Brain Behav Immun* 25:181–213.
- Tynan RJ, Naicker S, Hinwood M, Nalivaiko E, Buller KM, Pow DV, *et al.* (2010): Chronic stress alters the density and morphology of microglia in a subset of stress-responsive brain regions. *Brain Behav Immun* 24:1058–1068.
- Iwashyna TJ, Ely EW, Smith DM, Langa KM (2010): Long-term cognitive impairment and functional disability among survivors of severe sepsis. *JAMA* 304:1787–1794.
- Weaver JD, Huang MH, Albert M, Harris T, Rowe JW, Seeman TE (2002): Interleukin-6 and risk of cognitive decline: MacArthur studies of successful aging. *Neurology* 59:371–378.
- Perry VH, Cunningham C, Holmes C (2007): Systemic infections and inflammation affect chronic neurodegeneration. *Nat Rev Immunol* 7:161–167.
- Ericsson A, Liu C, Hart RP, Sawchenko PE (1995): Type 1 interleukin-1 receptor in the rat brain: Distribution, regulation, and relationship to sites of IL-1-induced cellular activation. *J Comp Neurol* 361:681–698.
- Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, Shen EH, Ng L, Miller JA, *et al.* (2012): An anatomically comprehensive atlas of the adult human brain transcriptome. *Nature* 489:391–399.
- Sukuki WA, Amaral DG (1994): Perirhinal and parahippocampal cortices of the macaque monkey: cortical afferents. *J Comp Neurol* 350:497–533.
- Critchley HD, Harrison NA (2013): Visceral influences on brain and behaviour. *Neuron* 77:624–638.
- Oitzl MS, van Oers H, Schobitz B, de Kloet ER (1993): Interleukin-1 beta, but not interleukin-6, impairs spatial navigation learning. *Brain Res* 613:160–163.
- Barrientos RM, Higgins EA, Sprunger DB, Watkins LR, Rudy JW, Maier SF (2002): Memory for context is impaired by a post context exposure injection of interleukin-1 beta into dorsal hippocampus. *Behav Brain Res* 134:291–298.
- Moore AH, Wu M, Shaftel SS, Graham KA, O'Banion MK (2009): Sustained expression of interleukin-1beta in mouse hippocampus impairs spatial memory. *Neuroscience* 164:1484–1495.
- Bellinger FP, Madamba S, Siggins GR (1993): Interleukin 1 beta inhibits synaptic strength and long-term potentiation in the rat CA1 hippocampus. *Brain Res* 628:227–234.
- Cunningham AJ, Murray CA, O'Neill LA, Lynch MA, O'Connor JJ (1996): Interleukin-1 beta (IL-1 beta) and tumour necrosis factor (TNF) inhibit long-term potentiation in the rat dentate gyrus in vitro. *Neurosci Lett* 203:17–20.
- Murray CA, Lynch MA (1998): Evidence that increased hippocampal expression of the cytokine interleukin-1 beta is a common trigger for age- and stress-induced impairments in long-term potentiation. *J Neurosci* 18:2974–2981.
- Schmid AW, Lynch MA, Herron CE (2009): The effects of IL-1 receptor antagonist on beta amyloid mediated depression of LTP in the rat CA1 in vivo. *Hippocampus* 19:670–676.
- Ban E, Haour F, Lenstra R (1992): Brain interleukin 1 gene expression induced by peripheral lipopolysaccharide administration. *Cytokine* 4:48–54.
- Gibertini M (1996): IL1 beta impairs relational but not procedural rodent learning in a water maze task. *Adv Exp Med Biol* 402:207–217.
- Pugh CR, Kumagawa K, Fleshner M, Watkins LR, Maier SF, Rudy JW, *et al.* (1998): Selective effects of peripheral lipopolysaccharide administration on contextual and auditory-cue fear conditioning. *Brain Behav Immun* 12:212–229.
- Banks WA, Erickson MA (2010): The blood-brain barrier and immune function and dysfunction. *Neurobiol Dis* 37:26–32.
- Rivest S (2009): Regulation of innate immune responses in the brain. *Nat Rev Immunol* 9:429–439.
- Saper CB, Romanovsky AA, Scammell TE (2012): Neural circuitry engaged by prostaglandins during the sickness syndrome. *Nat Neurosci* 15:1088–1095.
- Goehler LE, Gaykema RP, Hansen MK, Anderson K, Maier SF, Watkins LR (2000): Vagal immune-to-brain communication: A visceral chemosensory pathway. *Auton Neurosci* 85:49–59.
- Hansen MK, O'Connor KA, Goehler LE, Watkins LR, Maier SF (2000): The contribution of the vagus nerve in interleukin-1 β -induced fever is dependent on dose. *Am J Physiol Regul Integr Comp Physiol* 280:929–934.
- Wan W, Wetmore L, Sorensen CM, Greenberg AH, Nance DM (1994): Neural and biochemical mediators of endotoxin and stress-induced c-fos expression in the rat brain. *Brain Res Bull* 34:7–14.
- Harrison NA, Brydon L, Walker C, Gray MA, Steptoe A, Dolan RJ, *et al.* (2009): Neural origins of human sickness in interoceptive responses to inflammation. *Biol Psychiatry* 66:415–422.
- Hosoi T, Yasunobu O, Nomura Y (2000): Electrical stimulation of afferent vagus nerve induces IL-1b expression in the brain and activates HPA axis. *Am J Physiol* 279:141–147.
- Doeller CF, King JA, Burgess N (2008): Parallel striatal and hippocampal systems for landmarks and boundaries in spatial memory. *Proc Natl Acad Sci U S A* 105:5915–5920.
- Bohbot VD, Kalina M, Stepankova K, Spackova N, Petrides M, Nadel L (1998): Spatial memory deficits in patients with lesions to the right hippocampus and to the right parahippocampal cortex. *Neuropsychologia* 36:1217–1238.
- Hingorani AD, Cross J, Kharbanda RK, Mullen MJ, Bhagat K, Taylor M, *et al.* (2000): Acute systemic inflammation impairs endothelium-dependent dilatation in humans. *Circulation* 102:994–999.
- Insausti R, Juottonen K, Soininen H, Insausti AM, Partanen K, Vainio P, *et al.* (1998): MR volumetric analysis of the human entorhinal, perirhinal, and temporopolar cortices. *AJNR Am J Neuroradiol* 19:659–671.
- Slotnick SD, Moo LR, Segal JB, Hart J (2003): Distinct prefrontal cortex activity associated with item memory and source memory for visual shapes. *Brain Res Cogn Brain Res* 17:75–82.
- Gabrieli JDE, Corkin S, Mickel SF, Growdon JH (1993): Intact acquisition and long-term retention of mirror-tracing skill in Alzheimer's disease and in global amnesia. *Behav Neurosci* 107:899–910.
- Bohbot VD, Corkin S (2007): Posterior parahippocampal place learning in H.M. *Hippocampus* 17:863–872.
- Ploner CJ, Gaymard BM, Rivaud-Pechoux S, Baulac M, Clemenceau S, Samson S, *et al.* (2000): Lesions affecting the parahippocampal cortex yield spatial memory deficits in humans. *Cerebral Cortex* 10:1211–1216.
- Sommer T, Rose M, Glascher J, Wolbers T, Buchel C (2005): Dissociable contributions within the medial temporal lobe to encoding of object-location associations. *Learn Mem* 12:343–351.
- Malkova L, Mishkin M (2003): One-trial memory for object-place associations after separate lesions of the hippocampus and parahippocampal region in the monkey. *J Neurosci* 23:1956–1965.

42. Suzuki WL, Amaral DG (1994): Perirhinal and parahippocampal cortices of the macaque monkey: Cortical afferents. *J Comp Neurol* 350: 497–533.
43. Eacott MJ, Gaffan EA (2005): The roles of perirhinal cortex, postrhinal cortex, and the fornix in memory for objects, contexts, and events in the rat. *Q J Exp Psychol B* 58:202–217.
44. Buckley MJ, Gaffan D (1998): Perirhinal cortex ablation impairs visual object identification. *J Neurol* 18:2268–2275.
45. Zhu XO, Brown MW, Aggleton JP (1995): Neuronal signalling of information important to visual recognition memory in rat rhinal and neighbouring cortices. *Eur J Neurosci* 7:753–765.
46. Fahy EL, Riches IP, Brown MW (1993): Neuronal activity related to visual recognition memory: Long-term memory and the encoding of recency and familiarity information in the primate anterior and medial inferior temporal and rhinal cortex. *Exp Brain Res* 96:457–472.
47. Norman G, Eacott MJ (2004): Impaired object recognition with increasing levels of feature ambiguity in rats with perirhinal cortex lesions. *Behav Brain Res* 148:79–91.
48. Gaffan EA, Eacott MJ, Simpson EL (2000): Perirhinal ablation in rats selectively impairs object identification in a simultaneous visual comparison task. *Behav Neurol* 114:18–31.
49. Buffalo EA, Bellgowan PSF, Martin A (2006): Distinct roles for medial temporal lobe structures in memory for objects and their locations. *Learn Mem* 13:638–643.
50. Ikegaya Y, Delcroix I, Iwakura Y, Matsuki N, Nishiyama N (2003): Interleukin-1 β abrogates long-term depression of hippocampal CA1 synaptic transmission. *Synapse* 47:54–57.
51. Canevari L, Richter-Levin G, Bliss TV (1994): LTP in the dentate gyrus is associated with a persistent NMDA receptor-dependent enhancement of synaptosomal glutamate release. *Brain Res* 667:115–117.
52. Cunningham C, Sanderson DJ (2008): Malaise in the water maze: Untangling the effects of LPS and IL-1 β on learning and memory. *Brain Behav Immun* 22:1117–1127.
53. Brydon L, Harrison NA, Walker C, Steptoe A, Critchley HD (2008): Peripheral inflammation is associated with altered substantia nigra activity and psychomotor slowing in humans. *Biol Psychiatry* 63:1022–1029.
54. Reichenberg A, Yirmiya R, Schuld A, Kraus T, Haack M, Morag A, *et al.* (2001): Cytokine-associated emotional and cognitive disturbances in humans. *Arch Gen Psychiatry* 58:445–452.
55. Krabbe KS, Reichenberg A, Yirmiya R, Smed A, Pedersen BK, Bruunsgaard H (2005): Low-dose endotoxemia and human neuropsychological functions. *Brain Behav Immun* 19:453–460.
56. Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, *et al.* (2000): Inflammation and Alzheimer's disease. *Neurobiol Aging* 21:383–421.
57. Bird CM, Chan D, Hartley T, Pijnenburg YA, Rossor MN, Burgess N (2010): Topographical short-term memory differentiates Alzheimer's disease from frontotemporal lobar degeneration. *Hippocampus* 20: 1154–1169.
58. van Halteren-van Tilborg IADA, Scherder EJA, Hulstijn W (2007): Motor-skill learning in Alzheimer's disease: A review with an eye to the clinical practice. *Neuropsychol Rev* 17:203–212.
59. Marsland AL, Gianaros PJ, Abramowitch SM, Manuck SB, Hariri AR (2008): Interleukin-6 covaries inversely with hippocampal grey matter volume in middle-aged adults. *Biol Psychiatry* 64:484–490.