CLINICAL INNOVATION: Fair & Effective Incentives for New Uses of Established Drugs

The Physician & Pharmacist

Moderator: Simon Spinks Bristows LLP

Presenters and Panellists:

- **Dr Amitava Banerjee** UCL Farr Institute of Health Informatics
- Prof. Janis Shute Professor of Pharmacology, University of Portsmouth
- Prof. Graham Russell Emeritus Professor of Musculoskeletal Pharmacology, NDORMS, University of Oxford
- Prof. Jayne Lawrence Head of the Division of Pharmacy and Optometry at Manchester University

UCL



What the data tells us about missed opportunities- a physician's perspective

Dr Amitava Banerjee ма мрн ррыі гнеа гана гезс гвср Associate Professor in Clinical Data Science, UCL Honorary Consultant Cardiologist, UCLH and Barts Health @amibanerjee1 8th February 2018

UCL




Missed opportunities

- 1. Poor prediction from preclinical studies
- 2. Definitive evidence not available until phase 3 RCT
- 3. Not enough (primary or secondary) use of existing drugs
- 4. Too much focus on drugs alone or in isolation
- 5. The way we gather and use data

UCL




- 1. Poor prediction from preclinical studies
- 2. Definitive evidence not available until phase 3 RCT
- 3. Not enough (primary or secondary) use of existing drugs
- 4. Too much focus on drugs alone or in isolation
- 5. The way we gather and use data



UCL





- 1. Poor prediction from preclinical studies
- 2. Definitive evidence not available until phase 3 RCT
- 3. Not enough (primary or secondary) use of existing drugs
- 4. Too much focus on drugs alone or in isolation
- 5. The way we gather and use data







- 1. Poor prediction from preclinical studies
- 2. Definitive evidence not available until phase 3 RCT
- 3. Not enough (primary or secondary) use of existing drugs
- 4. Too much focus on drugs alone or in isolation
- 5. The way we gather and use data









<u></u> UCL



- 1. Poor prediction from preclinical studies
- 2. Definitive evidence not available until phase 3 RCT
- 3. Not enough (primary or secondary) use of existing drugs
- 4. Too much focus on drugs alone or in isolation
- 5. The way we gather and use data

