

UCL CERTIFICATE OF HIGHER EDUCATION IN ASTRONOMY

RELEVANT BASIC CONCEPTS OF MATHEMATICS (27th Feb 2020)

These guidelines give an idea of the basic concepts that potential applicants need to be familiar with, before the course starts in October 2020. Some of this material will be seen in detail during the course.

Rather than printing, we advice to use them to compile your own personal notes on a hand written dedicated notebook. This process helps fixing ideas much better.

The most relevant concepts of mathematics and physics are also illustrated in the boxes called TOOLS OF THE ASTRONOMER'S TRADE throughout your Universe text book.

best regards,

francisco diego

f.diego@ucl.ac.uk

NOTES ON MATHEMATICS

Mathematics provides a language in which we can express the relationships that arise in a quantitative area of science such as astronomy. With it, we can be both more concise and more precise. In theoretical branches of the subject, it is possible to work out formulae that help us to understand the underlying physics of a situation, such as the conditions in an astronomical object.

In the Certificate course we try to keep the use of mathematics to a minimum and you should be able to understand most of the material quite well even if you ‘bleep over’ equations when they arise. Where we do use equations, they are there to help you to understand the astronomy more fully.

In some of the homework, simple calculations are to be done, generally by substituting numbers into equations. We set these because it has been found that by getting hands-on experience of such calculations you will have a better feel for the physics and astronomy of what is going on. Some of the practical experiments also require simple calculation.

In the written examinations that come at the end of the year there may be a very few questions involving calculation. If you are comfortable in doing these they can give a quick way to accumulate good marks. If not, however, there is sufficient choice in the questions that the calculations can easily be avoided.

These notes are intended to give you a reminder of some of the basic properties of mathematical equations and how they can be used in a simple way. If you should have any difficulty with the maths in the course (or indeed with any aspect of it), then do please ask for help.

Introduction to Algebra

Algebra is a system of calculation where the numbers are represented by symbols. This has the advantages that general expressions can be written down, true for all numerical values of the symbols, and that manipulation can lead to other results—the calculation is done once for all rather than separately each time the numbers are changed.

Generally, each quantity is represented by one symbol, normally a letter of the alphabet a , b , etc, supplemented by Greek letters α , β , etc. [You will find a list of Greek letters and their names on the Certificate web site. These two are *alpha* and *beta*]. Both lower-case and upper-case letters can be used, and normally they have different meanings, so we must be careful to be consistent and always to write each letter the same way. It is wrong, for instance, to use upper-case M and lower-case m as though they are interchangeable, or R and r , and so on.

In some applications, particular two-letter combinations can be defined as a single quantity. For example, in calculus (a collection of techniques *not* required in the Certificate course) the combination dx is taken to represent a very small change in some quantity x . In the same way, sometimes δx or Δx [these are lower-case and upper-case Greek *delta*, respectively] might be used for a not-quite-so-small change in the value of x .

In physics, generally a symbol is used to represent some physical quantity. For example, d might represent a distance, r a radius or distance from a centre, A an area, and so on. But other letters could be used for any of these quantities, and these letters could be used elsewhere with other meanings. In any piece of algebra, the meanings attached to each symbol used should be made clear, either from the context or by actually defining them all.

In physics, some quantities can be considered as *variables*, such as a radial distance r , and others as *constants*. There are two types of constant, mathematical constants which are pure numbers and physical constants which have to do with the real world. An example of a mathematical constant is the ratio of the circumference of a circle to its diameter, conventionally written as π [lower case *pi*] and with a value 3.14159... that is given on any pocket calculator. Physical constants include all sorts of things that can be measured such as the mass of the Sun M_{\odot} or the gravitational constant G .

Combining quantities

Addition and Subtraction

If we want to add two quantities a and b we write the sum as $a + b$. If we want to subtract them, we write $a - b$.

For example, if $a = 4$ and $b = 3$ then

$$a + b = 4 + 3 = 7,$$

$$a - b = 4 - 3 = 1.$$

There can be more than two items in the list. For example, if we also have $c = 7$ and $d = 6$ then

$$a - b - c + d = 4 - 3 - 7 + 6 = 0.$$

In this case the result is zero. In other cases, some of the quantities or the final result could be negative numbers. So, we could have $e = -6$, $f = -2$ and

$$d - a + e - f = 6 - 4 - 6 - (-2) = -4 + 2 = -2.$$

Note that subtracting a negative number is equivalent to adding the corresponding positive number. Note also that if we have to multiply two minus signs it is clearer if we use brackets, writing $-(-2)$ rather than $--2$.

Multiplication

For numbers, the multiplication sign \times is used, as in $3 \times 4 = 12$. In algebra, if two symbols are written together, as ab , then they are to be multiplied. The sign \times also can be used, as $a \times b$, and so can a dot between the symbols, as $a \cdot b$. So, with $a = 4$ and $b = 3$ as before,

$$\begin{aligned} ab &= a \times b = a \cdot b \\ &= 3 \times 4 = 12. \end{aligned}$$

There is no rule about which way to write it in a particular situation, except that it's often best to keep things looking simple which favours the use of ab .

With numbers, the dot notation should not be used, to avoid confusion with a decimal point in an expression such as 2.5.

Division

To divide two numbers, we can use the division sign \div . Thus, $4 \div 2 = 2$. The sign \div can be used in algebra, as $a \div b$, but it is more common to use a line. In a display equation it can be written as

$$\frac{a}{b}$$

and to save space when writing a formula in the middle of text the forward slash or solidus $/$ can be used, as a/b . This is consistent with the ways in which we can write a fraction, $\frac{1}{2} = 1/2$.

Brackets

Brackets give a convenient way to write things more simply. For example, if we have the combination $bc + bd$ it is simpler and generally involves less writing to put this as

$$bc + bd = b(c + d).$$

In words, b times c plus b times d equals b times (c plus d).

We could have several sets of brackets, such as $(a + b)(c + d)$. Multiplying this out, we see that it equals

$$(a + b)(c + d) = ac + ad + bc + bd.$$

If we put in particular numerical values, then we get the same answer if we work out $a + b$ and $c + d$ and multiply them, or if we work out the four multiplications on the right-hand-side of this equation and add them up. It's quicker to do it the first way.

Brackets also are useful when we are dealing with fractions. Consider the expression

$$\frac{a+b}{c+d}$$

To write this with a slash, the correct form is $(a+b)/(c+d)$. It could be tempting to write $a+b/c+d$, but that actually means

$$a + \frac{b}{c} + d,$$

which is not at all the same.

This has to be kept in mind when using a calculator. Suppose we want to find $(2+3)/(4+6)$, which obviously is $5/10$ or $1/2 = 0.5$. If we key it in without brackets, we are likely to get 8.75 which is seriously wrong. If your calculator has brackets, use them in a case like this. Otherwise you could work out the numerator $2+3$ and denominator $4+6$ separately and then divide them.

In the fraction $(a+b)/(c+d)$, $(a+b)$ is called the *numerator* and $(c+d)$ the *denominator*.

We also have to be careful about the way we may split up a fraction like this. In fact,

$$\frac{a+b}{c+d} = \frac{a}{c+d} + \frac{b}{c+d} = a/(c+d) + b/(c+d).$$

There could be a temptation to think that it equals, for example, $a/c + b/d$ but this is not correct, as you may verify by putting in some numbers.

Powers

Integer Powers

If we multiply the same quantity by itself several times then we can express this as a power of that number. For instance,

$$a \times a = a^2,$$

$$a \times a \times a = a^3,$$

$$a \times a \times a \times a = a^4,$$

Powers are represented by exponents. For example 2, 3, 4 and (3-2) are all exponents.

and so on. Powers combine in a simple way,

$$a^2 \times a^3 = a^{(2+3)} = a^5,$$

or

$$a^3/a^2 = a^{(3-2)} = a^1 = a.$$

We read a^2 in words as 'a squared' and a^3 as 'a cubed'. For a^4 we say 'a to the power of 4' or 'a to the fourth power', and so on for other numbers.

As you see from this example, anything to the power of 1 equals itself. If we have an expression such as a^2/a^2 then this clearly must equal 1 (think of $a = 2$ and $2^2/2^2 = 4/4 = 1$). But also

$$a^2/a^2 = a^{(2-2)} = a^0$$

and in fact any number to the power of zero equals 1.

A special case is that one to any power still equals one, $1^n = 1$.

We can also have negative powers, such as

$$a^2/a^4 = a^{(2-4)} = a^{-2} = 1/a^2,$$

which is ‘one over a squared’ or ‘ a to the minus two’.

One over a number, or a number to the power of minus one, $1/a = a^{-1}$, is given a special name. It is the *reciprocal* of the number.

The power to the -1 gives another way to write a fraction. Thus

$$\frac{a}{b} = a/b = ab^{-1}.$$

This is often used with units, such as velocity. Kilometres per second can be written as km/s or as km s^{-1} .

Powers of 10

In Physics and Astronomy we often have to work with numbers that are very big or very small. To avoid writing a lot of figures or decimal places we make a great deal of use of powers of 10. So

$$10^0 = 1$$

$$10^1 = 10$$

$$10^2 = 100$$

$$10^3 = 1000$$

$$10^4 = 10000$$

and so on; and

$$10^{-1} = 0.1$$

$$10^{-2} = 0.01$$

$$10^{-3} = 0.001$$

$$10^{-4} = 0.0001$$

Fractional Powers and Roots

Suppose $b = a^2$ and we want to find an expression for a . We can do this by *taking the square root* of both sides of this relation, so that

$$a = \sqrt{b}.$$

The square root of b is a number which when multiplied by itself gives b and it is convenient to write this as $b^{1/2}$ so that

$$a^2 = (\sqrt{b})^2 = (b^{1/2})^2 = b^{1/2} \times b^{1/2} = b^{1/2+1/2} = b.$$

We can take other roots too,

$$\sqrt[3]{b} = b^{1/3},$$

$$\sqrt[4]{b} = b^{1/4},$$

and so on. We may write $b^{1/2}$ as $b^{0.5}$ and $b^{1/4}$ as $b^{0.25}$ but this doesn't work in every case—for example, $b^{1/3} = b^{0.333333\dots}$ which goes on for ever.

The reciprocal of a root is simply defined, for example

$$\frac{1}{\sqrt{x}} = \frac{1}{x^{1/2}} = x^{-1/2} = x^{-0.5}.$$

The integer and fractional powers can be combined in a variety of ways. For example, $a^{5/2}$ is the square root of a^5 . It is also the fifth power of the square root of a . So

$$a^{5/2} = \sqrt{(a^5)} = (a^5)^{\frac{1}{2}} = (\sqrt{a})^5 = (a^{1/2})^5.$$

Note that to take the power of a power the numbers are to be multiplied, $\frac{1}{2} \times 5 = \frac{5}{2}$. This quantity is *not* the same thing as $a^5 \times a^{1/2}$ which is actually $a^{(5+1/2)}$ or $a^{11/2}$.

Special Symbols

Equality

In an equation, $a = b$, the two sides have the same value. Mathematicians consider two different types of equation, those which are true whatever the values of the quantities involved, such as

$$a(b + c) = ab + ac,$$

and those which are true only for certain particular values, such as

$$x = a + b.$$

If $a = 2$ and $b = 3$ this is satisfied only by $x = 5$. The first case is called an *identity* and is sometimes written with \equiv instead of $=$,

$$a(b + c) \equiv ab + ac.$$

Sometimes we may want to say that two quantities are *approximately* the same. For instance, the number of seconds in a year can be worked out precisely but in some calculations it is enough to use the approximate value 3×10^7 . We could then write

$$1 \text{ year} \sim 3 \times 10^7 \text{ seconds.}$$

The symbols \sim , \approx and \simeq all can be used for approximate equality.

If two quantities are not equal we can write it as

$$2 + 2 \neq 5.$$

The symbol \propto is used to indicate that the left hand side is *proportional* to the right hand side, without giving the full relation or other factors that come in. For example, for a planet in orbit around the Sun the gravitational force is

$$F = \frac{GmM}{r^2}$$

where G is the gravitational constant and M and m are the two masses. If we want to emphasise the dependence on the distance r we could write

$$F \propto \frac{1}{r^2},$$

leaving out factors that are for the present unimportant.

To express inequality, we can use $>$ (greater than) or \geq (greater than or possibly equal to) or \gg (much greater than), with corresponding symbols $<$, \leq and \ll for less than.

Thus,

$$2 + 2 > 3,$$

$$1 \ll 10^9.$$

Summation and Multiplication

The Greek letter Σ (which is an upper case *sigma* and quite different in appearance from the lower case σ) is used to indicate the sum of a series of terms. Often the starting and finishing values of an index are added as subscript and superscript. So we have

$$\sum_{n=1}^4 n^2 = 1^2 + 2^2 + 3^2 + 4^2 = 1 + 4 + 9 + 16 = 30.$$

If there are n stars in a cluster, with individual masses m_i , the total mass M of the cluster is

$$M = m_1 + m_2 + m_3 + \dots + m_n = \sum_{i=1}^n m_i.$$

In the same way, Π which is upper case *pi* is used for a product:

$$\prod_{n=1}^4 n^2 = 1^2 \times 2^2 \times 3^2 \times 4^2 = 1 \times 4 \times 9 \times 16 = 576.$$

Working things out

Suppose we have a complicated expression such as

$$(a + b) \left(c - \frac{(d + k)}{e} \right) - (f - gj)(p^{h+k/l} + q)$$

where we are given numerical values for all the symbols. The correct order to work things out is

first, functions

next, expressions in brackets

next, multiplications and divisions

next, additions and subtractions

An example of a function here is taking a power of p . Brackets include the numerator or denominator of a fraction, or an exponent, even if brackets are not explicitly used in the formula. In a complicated expression, with brackets within brackets, we start with what is inside and work out.

Before we start, a word about terminology. In an expression like this, the quantities that are added or subtracted are called *terms*. In a product, the things to be multiplied are called *factors*. It's not uncommon for people to say *terms* when they mean *factors*.

In the first term here, we first work out the bracket $(d + k)$ and then the fraction $(d + k)/e$. We can then work out the two brackets $(a + b)$ and $(c - (d + k)/e)$ and multiply them together.

In the second term, we first find k/l , then $h + k/l$, and then work out the power $p^{h+k/l}$. We then add this to q to get the second factor. For the first factor, first multiply g by j , then subtract the result from f . The two factors can then be multiplied.

Finally, we can subtract the second term from the first one to get our answer.

Rearranging Equations

Very often we want to rearrange an equation to get an expression for one or other of the quantities involved. We can change an equation in a variety of ways, so long as we do exactly the same to both sides of the equation. We could:

- multiply or divide by any number or formula;
- add or subtract anything;
- take the reciprocal;
- take any power (e.g., square or take the square root);
- take any function (e.g., take 10 to the power of each side);
- exchange the two sides of the equation: if $a = b$ then $b = a$.

In a few cases we have to be careful as there can be multiple solutions. In particular, a is a square root of a^2 but so also is $-a$, because $(-a) \times (-a) = a \times a = a^2$. So for the square root of $b = a^2$ we have to say that $b^{1/2} = a$ or $-a$.

Consider the formula for the gravitational force between two objects of mass M and m a distance r apart:

$$F = \frac{GMm}{r^2}. \quad (\text{see chapter 4 of your book})$$

Suppose we want to find an expression for the mass M . We can first multiply both sides by r^2 :

$$F \times r^2 = \frac{GMm}{r^2} \times r^2 = GMm.$$

Next, divide through by Gm :

$$\frac{Fr^2}{Gm} = \frac{GMm}{Gm} = M,$$

as we can cancel out the G and the m in the numerator and denominator on the right

hand side. Finally, interchange the two sides to obtain

$$M = \frac{Fr^2}{Gm}.$$

Starting from the same formula, let us find an expression for r . We start by inverting both sides (that is, taking the reciprocal)

$$\frac{1}{F} = \frac{r^2}{GMm}$$

and exchanging the two sides

$$\frac{r^2}{GMm} = \frac{1}{F}.$$

Next, multiply both sides by GMm :

$$\frac{r^2}{GMm}GMm = \frac{1}{F}GMm$$

or

$$r^2 = \frac{GMm}{F}.$$

Then take the square root:

$$r = \pm\sqrt{\frac{GMm}{F}}$$

where \pm means *either plus or minus*. In this case we can reject the possible negative root, for the distance between two objects always is a positive number, so

$$r = \sqrt{\frac{GMm}{F}} = \left(\frac{GMm}{F}\right)^{\frac{1}{2}},$$

Logarithms

Definition: Logarithm of a number is the exponent or power to which another number called base has to be raised in order to get that number. The most common base number is 10, but there are others.

For example: the logarithm of 100 is 2 because 10 needs to be raised to the power 2 in order to get 100

Logarithms to base 10

The logarithm of a number is an example of what is called a *function*, which is a prescription for getting another number. In this case, we have two related equations,

$$a = \log b \text{ and } b = 10^a.$$

Logarithms satisfy these relations:

$$\log 1 = 0$$

$$\log 10 = 1$$

$$\log 100 = 2$$

$$\log 1000 = 3 \text{ etc}$$

$$\log(a^c) = c \log a$$

$$\log(ab) = \log a + \log b$$

$$\log\left(\frac{a}{b}\right) = \log a - \log b.$$

Logarithms are used in measuring stellar magnitudes. See chapter 19 of your book.

Exponential form is used to express extremely large or extremely small numbers

For example:

$$480\ 000\ 000 = 4.8 \times 10^8 = 4.8\text{E}+8$$

$$0.000\ 002\ 4 = 2.4 \times 10^{-6} = 2.4\text{E}-6$$

(the 'E' form is the same thing but easier to type. It is equivalent to the EXP key in calculators, which also have the +/- key useful to input negative exponents)

Multiply one by the other:

$$480\ 000\ 000 \times 0.000\ 002\ 4$$

$$\text{Same as: } 4.8 \times 2.4 \times 10^{8-6} = 11.52 \times 10^2 = 1152$$

$$\text{Same as: } 4.8 \times 2.4\text{E}+8-6 = 11.52\text{E}+2 = 1152$$

Now, division challenge! $480\ 000\ 000 / 0.000\ 002\ 4$

$$\text{Same as: } 4.8\text{E}+8 / 2.4\text{E}-6 = 2.0\text{E}8+6 = 2.0\text{E}+14$$

$$\text{Same as: } 200\ 000\ 000\ 000\ 000$$

Note E-6 in denominator climbs to numerator as E+6

TRIGONOMETRY

This is one of the most important parts of mathematics in our course, related to angular separations between objects in the sky and also their apparent sizes.

Trigonometry is based in the right angle triangle and the relative lengths of its sides related to the angles (in this case theta θ). a is called adjacent side, b is called opposite side and c is called hypotenuse.

There are three fundamental trigonometric functions:

$$\sin \theta = b/c \text{ (called sine of theta)}$$


$$\cos \theta = a/c \text{ (called cosine of theta)}$$

$$\tan \theta = b/a \text{ (called tangent of theta)}$$

These functions are available in any scientific calculator (i.e. smart phone these days).

Pythagoras theorem:

$$a^2 + b^2 = c^2$$




ANGULAR MEASUREMENTS

One of the main discoveries of ancient incipient scientists was that dividing the length of a circumference by its diameter, gave a number with infinite decimal places. That number is called Pi (π) and is equivalent to 3.141592653589 etc. For that reason Pi is called an irrational number.

Dividing the length of a circumference by the length of its radius, should give the value of the full-circle angle in radians. As the radius is half of the diameter, the number of radians in a circle is 2π or 6.28...rad. Conclusion: $360^\circ = 2\pi$ rad a very useful conversion factor to remember.

$$360 / 2\pi = 360/6.28\dots = 57.296^\circ \text{ which is the value of one radian in degrees}$$



Definition of **π**

$$\pi = L / d = 3.14159\dots$$

The length L is π times d
or π times $2 r$

ANGULAR UNITS

Astronomy deals with distant objects that we can not measure directly from Earth. The only size that can be measure is the angle that they appear to subtend in the sky.

The most common unit to measure angles is the DEGREE (symbol $^{\circ}$).


One degree results from dividing a full circle in 360 equal parts. 360 is an arbitrary number but very convenient as it can be divided by 2, 3, 4, 5, 6, 8, 9, 10, etc resulting in integer numbers.

One degree can be divided into 60 parts (again a convenient number) called arcminutes (arcmin), which in turn can be divided into 60 parts called arcseconds (arcsec). Therefore one degree has 60 arcmin or 3600 arcsec. For example, the constellation of Orion has an extent of 20° in the sky, while the Moon has only 0.5° or 30 arcmin.

There is another unit to measure angles which is far more practical for many astronomical applications, including telescopes and instruments. It is called RADIAN (rad).

One radian is the angle between two radii of a circle, where the arc segment between the two radii has the same length of the radius. In other words, we are dealing with a kind of equilateral triangle where one of the sides is an arc of a circle. Therefore, the radian is just short of 60° .

The value of an angle in radians is the ratio between the length of the arc within that angle to the length of the radius of the arc.



Definition of a radian

There are 2π radians in
a complete circumference

$$360^\circ = 2\pi \text{ rad}$$

$$\text{One rad} = 360^\circ / 2\pi = 57.29578^\circ$$


VERY SMALL ANGLES

Astronomy deals in general with very small, or narrow angles mainly because the objects to be measured are so far away. The Sun and the Moon as seen from Earth, appear so have the same angular diameter (by coincidence) of about 0.5° . Planets and distant stars show diameters far narrower than this, so the most common unit is the arcsecond, or $1/3600$ th of a degree. For example, the planet Jupiter has an angular diameter of about 40arcsec and the star Betelgeuse, which has the largest apparent diameter of all stars as seen from the Earth is only 0.05arcsec wide. As a useful reference, remember that a one pound coin will appear to have a diameter of one arcsec seen at a distance of nearly 5km! So one arcsec is a very small angle indeed.

$$\text{One rad} = 360^\circ / 2\pi = 57.29578^\circ$$

$$\text{Or } 57.29578 \times 60 \times 60 = 206265 \text{ arcsec}$$

One radian has 206265 arcsec



One arcsec is $1 / 206265$ rad

$$\text{Hence, } \tan (1 \text{ arcsec}) = 1 / 206265$$



22.5 mm

One arcsec



$$22.5 \text{ mm} \times 206265 = 4.64 \text{ E}+6 \text{ mm} = 4.64 \text{ km}$$

For example, a one pound coin would appear one arcsec wide when seen almost 5km away

IMPORTANT FORMULAE

Length L of a circumference of diameter d (or radius r)

$$L = \pi d = 2 \pi r$$

Surface area A of a circle of radius r

$$A = \pi r^2$$

Surface area S of a sphere of radius r

$$S = 4 \pi r^2$$

Volume V of a sphere of radius r

$$V = \frac{4}{3} \pi r^3$$