

A Hausdorff-Measure Boundary Element Method for Scattering by Fractal Screens II: Numerical Quadrature

Andrew Gibbs^{1,*}, David Hewett¹, Andrea Moiola²

¹Department of Mathematics, University College London, UK

²Dipartimento di Matematica, Università Delgi Studi di Pavia, Italy

*Email: andrew.gibbs@ucl.ac.uk

Abstract

In a related talk [1], the Boundary Element Method (BEM) is generalised to the case of scattering by fractal obstacles. Implementation requires evaluating integrals of singular Green’s kernels over fractal domains, with respect to *Hausdorff measure*. This motivated the development of new quadrature rules, which are discussed here.

Keywords: Quadrature, BEM, Fractals

1 Introduction

We will study numerical quadrature rules for the evaluation of integrals of the form

$$I_{\Gamma, \Gamma'}[\Phi] := \int_{\Gamma} \int_{\Gamma'} \Phi(x, y) d\mathcal{H}^d(y) d\mathcal{H}^d(x), \quad (1)$$

where Γ and Γ' are compact subsets of \mathbb{R}^2 of Hausdorff dimension $d > 0$ and $d' > 0$ respectively, \mathcal{H}^d and $\mathcal{H}^{d'}$ are the corresponding Hausdorff measures, and $\Phi(x, y) = \frac{e^{ik|x-y|}}{4\pi|x-y|}$ is the fundamental solution for the Helmholtz equation with wavenumber $k > 0$ in \mathbb{R}^3 . (In what follows, similar results hold for the analogous problem posed in \mathbb{R}^2 .)

Our motivation for approximating (1) is the *Hausdorff BEM*, which is introduced and analysed in the talk [1]. Such BEMs can model scattering by planar screens with non-integer (fractal) dimension, i.e. $d \in (1, 2)$.

2 Attractors of Iterated Function Systems

Now we describe in detail the class of fractal scatterers that we consider. An iterated function system (IFS) is a set of $2 \leq M \in \mathbb{N}$ *contracting similarities* $s_m(x) = \rho_m A_m x + \delta_m$, with contraction factors $\rho_m \in (0, 1)$, rotation matrices $A_m \in \mathbb{R}^{n \times n}$ and translations $\delta_m \in \mathbb{R}^n$, for $m = 1, \dots, M$. Saying that Γ is the *attractor* of the IFS means that Γ is the unique non-empty compact set satisfying $\Gamma = s(\Gamma)$, where $s(E) := \bigcup_{m=1}^M s_m(E)$, $E \subset \mathbb{R}^n$.

Our quadrature rules are based on splitting Γ into sub-components, using the IFS structure.




Figure 1: Vector indices on Cantor Dust.

To describe these sub-components we adopt vector index notation. For $\ell \in \mathbb{N}$ let $I_\ell := \{1, \dots, M\}^\ell$. Then for $E \subset \mathbb{R}^n$ let $E_0 := E$, and for $\mathbf{m} = (m_1, \dots, m_\ell) \in I_\ell$ define $E_{\mathbf{m}} := s_{\mathbf{m}}(E)$ and $s_{\mathbf{m}} := s_{m_1} \circ \dots \circ s_{m_\ell}$. For an illustration of this notation in the case of the middle-third Cantor dust see Figure 1. We say Γ is *hull-disjoint* if

$$\mathcal{R} := \min_{m \neq m'} \{\text{dist}(\text{Hull}(\Gamma_m), \text{Hull}(\Gamma_{m'}))\} > 0.$$

A key ingredient is the set of vector indices

$$L_h(\Gamma) := \{ \mathbf{m} = (m_1, \dots, m_\ell) \in \bigcup_{\ell \in \mathbb{N}} I_\ell : \text{diam}(\Gamma_{\mathbf{m}}) \leq h \text{ and } \text{diam}(\Gamma_{(m_1, \dots, m_{\ell-1})}) > h \}.$$

Heuristically, these indices correspond to a partition of Γ , where we have subdivided *just enough* so that all components have diameter no more than h . This is depicted in Figure 2.

3 The barycentre rule

We define the *barycentre rule* for double integrals:

$$Q_{\Gamma, \Gamma'}^h[f] := \sum_{\mathbf{m} \in L_h(\Gamma)} \sum_{\mathbf{m}' \in L_h(\Gamma')} w_{\mathbf{m}} w'_{\mathbf{m}'} f(x_{\mathbf{m}}, x'_{\mathbf{m}'}), \quad (2)$$




Figure 2: Partitioning Koch snowflake by $L_{0,3}(\Gamma)$. Barcentres $x_{\mathbf{m}}$ are represented by \times .

where the weights and nodes are given by $w_{\mathbf{m}} := \mathcal{H}^d(\Gamma_{\mathbf{m}})$ and $x_{\mathbf{m}} := \int_{\Gamma_{\mathbf{m}}} x \, d\mathcal{H}^d(x) / \mathcal{H}^d(\Gamma_{\mathbf{m}})$ for $\mathbf{m} \in L_h(\Gamma)$, with analogous definitions for Γ' . The weights and nodes can be easily computed in terms of the IFS parameters, see [2, (27-29)]. For the single integral version of (2), see [2, §3.1]. In all estimates that follow, C denotes a constant which depends only on Γ .

Theorem 1 (Lipschitz integrands) [2, Theorem 3.7] *If $L_0[f]$ and $L_1[f]$ are the Lipschitz constants of f and ∇f respectively in $\text{Hull}(\Gamma) \times \text{Hull}(\Gamma')$,*

$$\left| I_{\Gamma, \Gamma'}[f] - Q_{\Gamma, \Gamma'}^h[f] \right| \leq CL_p[f] h^{p+1} \text{ for } p \in \{0, 1\}.$$

A result for non-diagonal entries of Hausdorff BEM matrices follows immediately:

Corollary 2 (Smooth Galerkin integrals) [2, Proposition 5.2]

If $R := \text{dist}(\text{Hull}(\Gamma), \text{Hull}(\Gamma')) > 0$, then

$$\left| I_{\Gamma, \Gamma'}[\Phi] - Q_{\Gamma, \Gamma'}^h[\Phi] \right| \leq Ch^2 \frac{1 + (kR)^{n/2+1}}{R^{n+1}}.$$

4 Singular integrals of Laplace kernels

In Hausdorff BEM, the diagonal matrix elements correspond to (1) with $\Gamma = \Gamma'$. Because $|\Phi(x, y)| \rightarrow \infty$ as $|x - y| \rightarrow 0$, the rule (2) cannot be directly applied to (1) in this case. We will derive a new method for evaluating the singular (Laplace) component of (2), denoted $\Phi_0(x, y) := |x - y|^{-1}$. Then, to evaluate (1) with $\Gamma = \Gamma'$, we use a *singularity subtraction*

technique, by considering the Lipschitz continuous function $\Phi_* := \Phi - \Phi_0$, and splitting the integral as follows

$$I_{\Gamma, \Gamma}[\Phi] = I_{\Gamma, \Gamma}[\Phi_0] + I_{\Gamma, \Gamma}[\Phi_*], \quad (3)$$

and evaluating both components separately.

By exploiting the self-similarity of Γ , we can express $I_{\Gamma, \Gamma}[\Phi_0]$ as a linear function of $I_{\Gamma_m, \Gamma_m}[\Phi_0]$ for $m = 1, \dots, M$, which leads to

$$I_{\Gamma, \Gamma}[\Phi_0] = \frac{\sum_{m=1}^M \sum_{m' \neq m}^M I_{\Gamma_m, \Gamma_{m'}}[\Phi_0]}{1 - \sum_{m=1}^M \rho_m^{2d-1}}, \quad (4)$$

representing a singular integral as a linear combination of smooth integrals. The smooth integrals of (4) can be approximated using (2); we denote this approximation by $Q_{\Gamma, \Gamma, 0}^h$.

Theorem 3 (Singular Laplace-type integrals) [2, Corollary 4.7] *If Γ is Hull-disjoint, then*

$$\left| I_{\Gamma, \Gamma}[\Phi_0] - Q_{\Gamma, \Gamma, 0}^h \right| \leq Ch^2 \mathcal{R}^{-3} \left(1 - \sum_{m=1}^M \rho_m^{2d-1} \right)^{-1}.$$

5 Approximating (1)

Noting the decomposition (3), Theorem 3 states that $I_{\Gamma, \Gamma}[\Phi_0]$ can be estimated with $O(h^2)$ error, provided Γ is hull-disjoint.

Since $\Phi_* \in C^{0,1}(\mathbb{R}^n) \setminus C^{1,1}(\mathbb{R}^n)$, Theorem 1 suggests $|I_{\Gamma, \Gamma}[\Phi_*] - Q_{\Gamma, \Gamma}^h[\Phi_*]| = O(h)$. With further work it can be shown that this is actually $O(h^2)$, when (i) Γ is hull-disjoint and (ii) $\rho_1 = \dots = \rho_M$. Hence using (3) we can approximate $I_{\Gamma, \Gamma}[\Phi]$ with $O(h^2)$ accuracy (see [2, §5] for details).

Furthermore, numerical experiments [2, §6] suggest $O(h^2)$ convergence for fractals which violate either or both of the conditions (i)–(ii).

References

- [1] S. N. Chandler-Wilde, A. Caetano, A. Gibbs, D. P. Hewett and A. Moiola, A Hausdorff-Measure Boundary Element Method for Scattering by Fractal Screens I: Numerical Analysis, in *Proceedings of the 15th International Conference on Mathematical and Numerical Aspects of Wave Propagation, Paris, France, 25-29 July 2022*.
- [2] A. Gibbs, D. P. Hewett and A. Moiola, Numerical quadrature for singular integrals on fractals, *arXiv* **2112.11793** (2021).