



Contents lists available at ScienceDirect

Journal of Quantitative Spectroscopy & Radiative Transfer

journal homepage: www.elsevier.com/locate/jqsrtCross-sections for heavy atmospheres: H₂O self-broadeningLara O. Anisman^{a,*}, Katy L. Chubb^b, Quentin Changeat^a, Billy Edwards^{c,a},
Sergei N. Yurchenko^a, Jonathan Tennyson^a, Giovanna Tinetti^a^a Department of Physics and Astronomy, University College London, Gower Street, London, WC1E 6BT, United Kingdom^b Centre for Exoplanet Science, University of St Andrews, North Haugh, St Andrews, KY16 9SS, United Kingdom^c AIM, CEA, CNRS, Université Paris-Saclay, Université de Paris, Gif-sur-Yvette, F-91191, France

ARTICLE INFO

Article history:

Received 24 September 2021

Revised 27 February 2022

Accepted 28 February 2022

Available online 4 March 2022

Keywords:

Exoplanet atmospheres

Water vapor

Opacities

Radiative transfer

Line broadening

Super-Earths

mini-Neptunes

ABSTRACT

The discovery of super-Earth and mini-Neptune exoplanets means that atmospheric signals from low-mass, temperate exoplanets are being increasingly studied. The signal acquired as the planet transits its host star, known as the *transit depth*, is smaller for these planets and, as such, more difficult to analyze. The launch of the space telescopes *James Webb* (JWST) & *Ariel* will give rise to an explosion in the quality and quantity of spectroscopic data available for an unprecedented number of exoplanets in our galaxy. Accurately extracting the information content, thereby permitting atmospheric science, of such data-sets will require robust models and techniques. We present here the analysis of simulated transmission spectra for water-rich atmospheres, giving evidence for non-negligible differences in simulated transit depths when self-broadening of H₂O is correctly accounted for, compared with the currently typically accepted standard of using H₂ and He-broadened cross-sections. Our case-study analysis is carried out on two super-Earths, focusing on water-based atmospheres, ranging from H₂-rich to H₂O-rich. The transit depth is considerably affected, increasing values by up to 60 ppm, which is shown to be detectable with JWST and *Ariel*. The differences are most pronounced for the lighter (i.e. $\mu \sim 4$) atmospheres. Our work illustrates that it is imperative that the field of exoplanet spectroscopy moves toward adapted cross-sections, increasingly optimized for high- μ atmospheres for studies of super-Earths and mini-Neptunes.

Crown Copyright © 2022 Published by Elsevier Ltd. All rights reserved.

1. Introduction

The field of exoplanet atmospheric spectroscopy relies heavily upon accurately derived cross-sections, generated for particular pressure and temperature ranges as well as for specific molecules. As the number of known exoplanets continues to increase rapidly, we are witnessing an influx of small, more temperate worlds ($\leq 10 M_{\oplus}$, ≤ 5000 K). The range of super-Earths and mini-Neptunes discovered has opened up the field of possibilities for observable atmospheres on these planets. Prime examples include the TRAPPIST-1 system, [1], 55 Cnc e [2,3], GJ 1132 b [4,5], GJ 1214 b [6], K2-18 b [7] and LHS 1140 b [8].

Small planets of increasing interest in the field are those that lie within the radius valley [9–11], i.e. between $1.5 - 2.0 R_{\oplus}$, whereby the dearth of planets in this region is theorized to be consistent with the intersection between super-Earths and sub-Neptunes or water-worlds. Planets smaller than $1.8 R_{\oplus}$ could have thinner, possibly H₂-depleted atmospheres (e.g. [12–14]), whilst planets with

radii larger than this threshold could possess volatile-rich atmospheres, H₂/He or H₂O-rich in particular (e.g. [15,16]). However, observational constraints, in the form of atmospheric transmission spectra, are needed to confirm these hypotheses and therefore the nature of these transitional planets. Furthermore, this population of small planets deviates from our understanding of planet atmospheres which has been mostly built upon our knowledge of hot-Jupiters and the solar system. For example, cross-sections that are utilized in the exoplanet field (based on the line lists provided by ExoMol [17], HITRAN [18], MoLLIST [19] and HITEMP [20]) are typically generated for atmospheres which are either dominated by H₂ and He, or air for Earth-like planets, in the case of HITRAN (see recent works, such as [21–24]). Whilst this is appropriate for gas giant planets or Earth-like planets, such cross-sections do not include effects such as the self-broadening of heavy molecules like H₂O. When considering atmospheres heavier than H₂/He, such as is expected to be prevalent on super-Earths and water worlds [16,25–28], this provides a non-negligible difference in the simulated (wavelength-dependent) atmospheric signal obtained during transit, known as the *transit depth*.

H₂ and He are light molecular species which only interact weakly with other molecules at long range and are therefore very

* Corresponding author.

E-mail address: lara.anisman.18@ucl.ac.uk (L.O. Anisman).

inefficient line broadeners. Water, conversely, is both heavier and possesses a significant dipole which leads to enhanced long-range interactions. Broadening by water vapour is known to be significant in the Earth's atmosphere even though it is generally only a trace species. As transit spectroscopy probes regions of an exoplanet atmosphere as it approach optical thickness, the increased line broadening by water can be expected to lead to significantly enhanced overall absorption compared to the case where only H₂ and He are considered.

Previous works have investigated the effects of various choices when computing cross-sections for modelling exoplanet atmospheres, such as [29–32]. These investigations included the choice of broadening parameters, although these typically focus on H₂/He atmospheres and associated broadening parameters. Recently there has been some work more focused on cross-sections for heavier atmospheres, such as [28,33].

In this work we investigate the effects of including H₂O self-broadening, in addition to H₂/He-broadening, in the calculation of our water cross-sections which we use to model exoplanet atmospheres. We find that the transit depth is considerably affected; the largest difference being found for one of our case-study planets GJ 1214 b, a 6.26 M_⊕, 2.85 R_⊕ super-Earth.

2. Methodology

2.1. Transmission spectroscopy

If the orbital plane of a planet around its host star is aligned approximately parallel to our line of sight with the system (analogous to 90° inclination), the planet will transit in front of its star. Assuming that there is no atmosphere, and that the planet is totally opaque to its incoming starlight, this transiting motion will cause a drop in the amount of stellar flux we receive from the host star. This change in detected light is known as the *transit depth*, ΔF , which is equal to the ratio between the surface area of the planet (as we view it, in 2D) and the surface area of the star. Since we assume both objects to be totally symmetrical, this reduces to:

$$\Delta F = \frac{F_{out} - F_{in}}{F_{out}} = \left(\frac{R_p}{R_*}\right)^2 \quad (1)$$

which gives a measure of the relative change in flux as the planet blocks its starlight. This provides us with an observable quantity with which we can quantify the size of the planet, if we know the stellar properties, which can be derived from models.

Now, if the planet possesses an atmosphere, an envelope of gas which surrounds the planet which is maintained by the planet's gravitational force, the molecules present will absorb, scatter and reflect incoming starlight, in addition to thermally emitting photons. Owing to the varied and distinct spectral characteristics of different molecules, how opaque a certain atmosphere is to incoming stellar flux will vary significantly with wavelength. This information is described by the quantity $\tau(\lambda)$, given in Eq. (2), known as the *optical depth*. Overall, regarding transmission spectroscopy, we can treat the atmosphere as a purely absorbing and single-scattering medium as a good (first-order) approximation for radiative transfer through the planetary atmosphere.

Given an arbitrary path through the atmosphere for which radiation transmits with wavelength-dependent initial intensity I_λ , the transmitted radiance will be attenuated by absorption and scattering processes. We can denote this reduction in intensity as a function of path ds as $dI_\lambda/ds = -I_\lambda\sigma_\lambda\rho$, where σ_λ is the total mass extinction cross section (the sum of the absorption and scattering cross sections) and ρ is the density of the medium. Integrating up and using the fact that the optical depth as a function of atmospheric height is determined by summing the opacity contribu-

tions of all molecular species present, we recover the *Beer-Bouguer-Lambert Law*:

$$I_\lambda(z) = I_\lambda(0)e^{-\tau_\lambda(z)} \quad \text{with} \quad \tau_\lambda(z) = \sum_m \int_z^{z_\infty} \sigma_{m,\lambda}(z') \chi_m(z') \rho(z') dz', \quad (2)$$

where χ_m and ρ are the column density of a given molecular species and the number density of the atmosphere, respectively.

We may now rewrite Eq. (1) as:

$$\Delta F = \frac{F_{out} - F_{in}}{F_{out}} = \left(\frac{R_p + h_\lambda}{R_*}\right)^2 \approx \frac{R_p^2 + 2R_p h_\lambda}{R_*^2} \quad O(h_\lambda) \quad (3)$$

where we may describe the atmospheric height function as:

$$2R_p h_\lambda = 2 \int_0^{z_{max}} (R_p + z)(1 - e^{-\tau_\lambda(z)}) dz, \quad (4)$$

where z_{max} denotes the height of the atmosphere.

Using this formalism, the transit depth of an atmosphere-bearing planet for any given wavelength may be calculated, using derived *cross-sections* (temperature, pressure, and wavelength dependent) for a given molecular species. If we populate a model atmosphere with a given temperature profile, pressure profile, chemical species abundances and a specified cloud distribution we may generate a transmission spectrum (ΔF vs. λ) for the atmosphere, thereby *forward-modelling* it. To date, there is extensive literature pertaining to the collection of transmission data, alongside analysis of the generated transmission spectra, for a wide variety of exoplanets; from hot-Jupiters [34–36] to habitable-zone super-Earths [1,7,8].

2.2. Broadening parameters

The Voigt profile is commonly used to represent line broadening in exoplanet atmospheres, which is a convolution of the temperature-dependent Gaussian line profile and the pressure-dependent (and therefore dependent on broadening species) Lorentzian profile. The equation for the Lorentzian line width (HWHM) for a given pressure P and temperature T , is given by:

$$\gamma_L = \gamma \left(\frac{T_0}{T}\right)^n \frac{P}{P_0}. \quad (5)$$

Here, T_0 and P_0 are the reference temperature and pressure, whilst γ and n are the reference HWHM and temperature exponent, respectively. The latter two terms are known as pressure-broadening parameters, and are dependent primarily on the molecular species being broadened and the species inducing it. Therefore these are the parameters which, upon variation, enable us to study the self-broadening effects of water considered in this study; considering atmospheres comprising varying levels of H₂O with respect to H₂ and He. The self-broadened H₂O cross-sections used in this study were computed using ExoCross [37], as were the H₂/He-broadened cross-sections for H₂O. The latter are similar to those from the ExoMolOP database [21] but with J -dependent broadening parameters (where J is the rotational angular momentum quantum number) from the ExoMol website¹ used for H₂ and He broadening [38,39]. All H₂O cross-sections presented here use the ExoMol POKAZATEL line list [40]. We computed the line wings out to 500 Voigt widths in all cases, out to a maximum of 25 cm⁻¹. The values of γ and n for the self-broadening of H₂O used in the present study are detailed below.

¹ https://www.exomol.com/data/data-types/broadening_coefficients/H2O/.

2.2.1. Self-broadened half-width, $\gamma_{\text{H}_2\text{O}}$

There are many literature sources with broadening parameters for self-broadening of H_2O . For example, Gamache and Hartmann [41] compiled and compared various parameters related to H_2O line shape, including values for the half-width γ for self- H_2O broadening. There are over 47,000 lines in their database, with values of $\gamma_{\text{H}_2\text{O}}$ ranging from 0.108 - 0.805 $\text{cm}^{-1} \text{atm}^{-1}$. The 2020 release of the HITRAN database [42], who follow a “diet” procedure [43], provide an update of this 2004 broadening measurement database of [41]. A simple average of all values of the main isotopologue of H_2O (with no weighting) from HITRAN2020 for $\gamma_{\text{H}_2\text{O}}$ yields a value of 0.35 $\text{cm}^{-1} \text{atm}^{-1}$. There are many other works with available broadening values: for example, [44] and [45] both present a number of values for $\gamma_{\text{H}_2\text{O}}$ for a few thousand lines each. They report average values of $\gamma_{\text{H}_2\text{O}} = 0.4$ and between 0.1 and 0.5 $\text{cm}^{-1} \text{atm}^{-1}$, respectively.

As noted above, for the present study we extract the broadening parameters from the HITRAN2020 [42] database as a function of rotational angular momentum quantum number J , computing an average value of γ for each value of J . The data extends up to a maximum of $J=26$, and vary between 0.1 $\text{cm}^{-1} \text{atm}^{-1}$ for high J to 0.5 $\text{cm}^{-1} \text{atm}^{-1}$ for low J .

2.2.2. Temperature exponent $n_{\text{H}_2\text{O}}$ for self-broadened half-width

HITRAN2020 [42] currently only include γ_{self} and not n_{self} values for H_2O , due to the large effort required to validate and populate such parameters into their database. For now we use averaged J -dependent values for γ_{self} from HITRAN2020 [42], and an apparently typical value of $n_{\text{self}}=0.7$ for H_2O . The focus of some ongoing and future work is to update the ExoMol [46] and ExoMolOP [21] databases to include broadening parameters in a more comprehensive way. This is not a simple undertaking: the H_2O line list used in this work, for example, ExoMol POKAZATEL [40], contains 6 billion transitions between 800,000 energy levels, with even larger line lists required to describe larger species, see Table 13 of [21].

Although it is beyond the scope of the current study to perform a comprehensive assessment of available temperature exponents for self-broadening of H_2O , there are a number of studies in the literature which have focused on analysing water vapor spectra at various temperatures in order to determine the temperature dependence n of the self-broadened half-width γ . Here, we summarize the results from a selection of works, but note that this is not a comprehensive sample. Grossmann and Browell [47] analyzed spectra of water vapor in the 720 nm region, finding an average value of $n=0.75$. Studies such as [48–50] all analyzed particular rotational lines in the low-wavenumber region of the spectra, between around 250 - 390 K. They find values of n_{self} of 0.62, 0.89, and 0.85, respectively. Both [49,50] look at various broadeners, including self. They find that both γ and n are generally larger for the cases where H_2O is the broadener, in comparison to N_2 , O_2 or Ar as broadeners (n is 0.52, 0.64, 0.49 for those cases, respectively for example in [49]). Alder-Golden et al. [51] analyzed low- J lines of H_2O close to 12,200 cm^{-1} in the 330 - 540 K temperature range. They find an average value of γ_{self} at 296 K of 0.456 $\text{cm}^{-1} \text{atm}^{-1}$ (compared to 0.095 $\text{cm}^{-1} \text{atm}^{-1}$ for air-broadening). The temperature coefficient n for self-broadening of H_2O was found to be 0.9 on average across the spectral region measured. Podobedov et al. [52] analyzed several lines of H_2O in the region around 12 - 52 cm^{-1} for temperatures between 263 - 340 K and over a pressure range from 0.0003 - 0.014 bar. The J - and T -dependent values of γ were found to vary between 0.67 - 1.07 $\text{cm}^{-1} \text{atm}^{-1}$. They found J -dependent values of n_{self} between 0.56 and 0.81.

Table 2 gives some average values extracted from the HITRAN 2020 [42] database for various molecules and broadeners. The H_2 - and He-broadened average values used in the ExoMolOP database

Table 1

a) stellar and planetary parameters for two small planets, for input into TauREx 3.0, derived from [61], [62], b) list of the forward-modeled parameters, their values and the scaling used.

a) stellar & planetary parameters			
parameter	GJ 1132 b	GJ 1214 b	
T_* [K]	3270	3026	
R_* [R_\odot]	0.207	0.220	
M_* [M_\odot]	0.181	0.176	
M_p [M_\oplus]	1.66	6.26	
R_p [R_\oplus]	1.13	2.85	
P_{orbital} [days]	1.63	1.58	
b) forward model parameters			
parameter	GJ 1132 b	GJ 1214 b	type
P_{clouds}	None	None	opaque
T [K]	500	600	iso.
$\log_{10} V_{\text{H}_2\text{O}} \in$	[-2, 2]	[-2, 2]	fill
V_{He}	$1e-7$	$1e-7$	trace

Table 2

Average values of γ and n for various species where parameters are available from HITRAN-2020 [42]. The custom data search from HITRANonline as described in [79] was used in order to extract the relevant parameters. Where ExoMolOP [21] is labelled as the source (where only H_2 and He-broadening was considered), the citations used to find the averaged values are listed in the footnote to this table.

Species	Broadener	γ	n	Source
CO_2	H_2O	0.14	0.79	HITRAN
	Self	0.09	0.64	HITRAN
	Air	0.07	0.71	HITRAN
	H_2	0.11	0.58	HITRAN
	He	0.06	0.3	HITRAN
	H_2	0.09	0.59	ExoMolOP
CO	H_2O	0.09	0.68	HITRAN
	Self	0.06	-	HITRAN
	Air	0.05	0.7	HITRAN
	CO_2	0.06	0.66	HITRAN
	H_2	0.07	0.58	HITRAN
	He	0.05	0.54	HITRAN
CH_4	H_2O	0.07	0.85	HITRAN
	Self	0.07	-	HITRAN
	Air	0.05	0.67	HITRAN
	H_2	0.06	0.6	ExoMolOP
	He	0.03	0.3	ExoMolOP
	Self	0.35	-	HITRAN
H_2O	Self	0.35	-	HITRAN
	Air	0.07	0.62	HITRAN
	H_2	0.06	0.2	ExoMolOP
	He	0.01	0.13	ExoMolOP

CH_4 : [80–88] CO : [89–94] H_2O : [38,95–102] CO_2 assumed parameters of C_2H_2 : [103]

[21] (a database tailored for modelling “hot Jupiter”-type exoplanet atmospheres) are also included for reference. It can be seen that in general that the largest γ and n values occur when H_2O is the broadener, which for $\gamma_{\text{H}_2\text{O}}$ is an order of magnitude larger than H_2 - and He-broadening. The contrast between H_2 - or He-broadening and H_2O -broadening, however, does not appear to be so large for other species as for water vapor. A similar observation was noted by [28], who highlight that self-broadening for H_2O is typically up to 7 times larger than H_2/He broadening based on their compilation of literature values.

We note that the power law for temperature dependence which we assume in this work may not work well over large temperature ranges, including those temperatures of the atmospheres we are




Fig. 1. Cross-sections computed using the self-broadening parameters for H₂O, for the abundances of water vapor given in the legend. We compare the self-broadened cross-sections (orange, pink) with the H₂/He-only broadened cross-sections (blue). Left: atmospheric pressure of 1 bar and right: atmospheric pressure of 10⁻¹ bar. In an atmosphere with 10% H₂O, we assume the remaining 90% atmosphere is comprised of H₂/He in solar abundances, and that the broadening is therefore 90% dominated by H₂/He and the remaining 10% from H₂O self-broadening.




Fig. 2. The contribution function, $\frac{d\tau}{dP}$, is defined as the wavelength-averaged variation in the optical depth, τ with pressure, P . On the LHS, τ is plotted in $\mu - P$ space; for which the contribution $\frac{d\tau}{dP}$ is normalized and displayed in the central panel; whilst on the RHS this contribution is plotted for varying water vapor abundance; illustrating that deeper pressures can be probed in lighter atmospheres.

modelling. This has been demonstrated by works such as [53], who developed the more advanced Gamache–Vispoel double power law (DPL) model. Other works such as [54] highlight non-Voigt effects to line broadening. We will consider updating our cross-sections using these more robust line-shape models in the future. We also expect to have more robust values for the self-broadening parameters of H₂O, and other molecular species, as a result of on-

going work from projects such as ExoMol [46] and HITRAN [55], and others [56,57], which we would like to incorporate into our cross-sections in the future. In Fig. 1 we present our derived H₂O-broadened cross-sections at $T = 600$ K for various pressures, in comparison to the standard H₂/He-broadened ones; in both cases it is clear that H₂O self-broadening widens the profiles of each absorption peak.




Fig. 3. Individual over-plots of the (resolution 200) spectra produced with both H_2 and H_2O -broadened cross-sections for each planet, for varying water abundance, with absolute differences beneath. Top: GJ 1132 b (500 K), bottom: GJ 1214 b (600 K).

2.3. Simulating transmission spectra

Equation (5) exhibits a linear relationship for γ_L with atmospheric pressure, P . Hence, we anticipate the strongest broadening effect to contribute deeper in the atmospheres we wish to model. Consequently, the abundance of water vapor in the atmosphere should affect our results, since although a decrease in molecule number density ought to minimize the overall opacity contribution induced by self-broadening, lighter atmospheres with low-mean-molecular weight allow us to probe deeper pressures; this is illustrated in Fig. 2, where the contribution function, $\frac{d\tau}{dP}$, is defined as the wavelength-averaged variation in the optical depth, τ with pressure, P . This informs our implementation of atmospheric models endowed with a variety of water vapor abundances, in order to examine for which abundances the self-broadening effects of H_2O are both most prominent, and, most observable with future space missions.

In order to simulate forward models of transmission spectra for GJ 1132 b and GJ 1214 b, this analysis was performed using the publicly available retrieval suite TauREx 3.0 [58–60]. For the stellar parameters and the planet mass, we used the values from [61] and [62] as given in Table 1 a). In our runs we assumed that the planets possess a range of different water-based atmospheres, with fill gas abundance ratio given by $V_{\text{H}_2\text{O}} = x$, with the

chosen values of x given in Table 1 b), and the remainder of the molecular abundance made up of H_2 and He, as specified. Additionally, we included the collision induced absorption (CIA) from $\text{H}_2\text{-H}_2$ [63,64] and $\text{H}_2\text{-He}$ [65], as well as Rayleigh scattering for all molecules. Finally, our simulated atmospheres are cloud-free, isothermal and have molecular abundance profiles which are constant with altitude. The assumptions of isothermal and isochemical atmospheres hold for interpreting current data [8,66]. While these approximations might be too simplistic to interpret accurately JWST and Ariel data [67,68], they will not change the conclusions of our paper. For each of the two planets two sets of spectra were generated: one using the standard H_2/He -broadened cross-sections, and another with the H_2O -broadened opacities, as described in Section 2.2.

3. Results

3.1. Forward-modelling of transmission spectra

We generate transmission spectra at a native resolution of 15,000 for all three planets, before binning down to a nominal resolution and signal-to-noise ratio of 200 and 10, respectively, with TauREx 3.0 (enabling eventual comparison with Ariel and JWST errorbars). We utilize both the H_2/He - and the H_2O -




Fig. 4. Comparison of simulated error bars for observation of a) GJ 1132 b, and b) GJ 1214 b with future-missions Ariel and JWST with the absolute wavelength-dependent differences in transit depth on the simulated transmission spectra, using H_2 - vs. H_2O -broadened cross-sections. In both cases these broadening-induced differences should be directly observable in the near-future.

broadened cross-sections described in Section 2.1, yielding the spectra given in Fig. 3, with planetary and stellar parameters described in detail in Table 1. In comparison with spectra produced using H_2/He -broadened cross-sections, it is evident that by using cross-sections calculated for the water-dominated atmospheres the spectral features are amplified due to the additional absorption achieved by including self-broadening of H_2O . In order to quantify these wavelength-dependent differences, these sets of spectra which are over-plotted the top panels of Fig. 3 have been subtracted to obtain the results presented in the lower panels, from which we determine maximum absolute differences in transit depth for the lightest secondary-type atmospheres (10% H_2O); of 60 and 20 ppm for GJ 1214 b and GJ 1132 b, respectively.

3.2. Ariel & JWST error-bars

During its primary mission, Ariel will survey the atmospheres of 1000 exoplanets [69,70] and many of these targets could be in the Super-Earth and Sub-Neptune regime [71]. Meanwhile, during the Guaranteed Time Observations (GTO) and first cycle of the General Observing (GO), around 70 planets will be observed with JWST [72]. Nearly half of these JWST targets have a radius of

$\leq 2.5R_{\oplus}$ and, therefore, may not have a H_2 -dominated atmosphere. During its lifetime, JWST is expected to observe a couple of hundred exoplanets [e.g. 73].

To investigate the detectability of broadening-induced differences with future instruments, we generated error bars for the simulated spectra. We then compared the size of these uncertainties to the absolute differences between the H_2 - and H_2O -broadened spectra. For Ariel, we generated error bars using ArielRad [74] while, for JWST, we used a modified version of the radiometric tool described in Edwards and Stotesbury [75] which utilizes the JWST instrument parameters from Pandeia [76].

For JWST we modeled observations with NIRISS GR700XD (0.8 - 2.8 μm) and NIRSpec G395M (2.9 - 5.3 μm), whilst for Ariel, which provides simultaneous coverage from 0.5 - 7.8 μm , we simulated error bars at tier 2 resolution. Presented in Fig. 4, for GJ 1132 b and GJ 1214 b, we observe that both instruments will be sensitive enough to reveal such differences when integrating multiple transits, specifically 45 and 40 in the case of Ariel (tier 2), and 10 and 5 in the case of JWST for GJ 1132 b and GJ 1214 b, respectively. We note that both these planets will be studied by JWST in the first cycle of observations.

4. Discussion

In all spectra, an increase in μ , corresponding to increased volume mixing ratio of water, corresponds to reduced scale height, since $H = \frac{kT}{\mu g}$ (where μ is the mean molecular weight of the atmosphere, g is the planet gravity and k is the Boltzmann constant). For the individual atmospheres we observe a saturation of features as the atmospheric mean-molecular-weight gets larger. In tandem, an increase in mean-molecular weight of the atmosphere results in a decrease in the atmospheric pressure at which we can probe, as illustrated in Fig. 2. Thus, the largest observed differences in transit depth vs. wavelength are found in the lightest secondary-type atmospheres, namely those with 10% water vapor, for all employed temperatures. As for the two planets considered, the largest transit depth increases are observed in GJ 1214 b, with an equilibrium temperature of 600 K, as opposed to HAT-P-11 b, which possesses an equilibrium temperature of 900 K.

Although the Lorentzian line profile for H₂O self-broadening is approximately inversely proportional to the atmospheric temperature, low-temperature planets have smaller global transit depths, due to the reduction in scale height the temperature induces. Hence, there is a trade-off between the strength of the broadening effect and detectability of such effects. Our work finds that the atmospheric temperatures for such differences to become detectable with future missions, as discussed, is thus ~ 600 K.

Thus, consideration and utilization of self-broadening effects induced by not just H₂O, but all large- μ molecules should have a stronger impact on the medium-cool atmospheres of temperate planets, as well as those with low-mean-molecular-weight secondary atmospheres, namely those closer to the threshold abundance of around 10% non-H₂ or -He which we define to the boundary between primary and secondary type atmospheres; precisely those planets which sit in the transition region between super-Earths and sub-Neptunes, whose atmospheres remain illusive and out-of-reach with current telescopes, due to known limits on both the signal-to-noise and resolution [8]. The next generation of telescopes will widen and deepen our spectral view, both in terms of signal-to-noise and resolution, but also in wavelength coverage; hence these problems should be easily tackled with Ariel and JWST. It is therefore especially imperative for the accuracy of cross-section data to compete with the level of precision obtainable with these future facilities.

Standard H₂/He-broadened cross-sections which are in widespread use by the exoplanet community are simply non-optimal for the study of secondary-type, heavy atmospheres dominated by non-H₂/He species due to the fact that these opacities are calculated with respect to a nominal atmosphere which is dominated by H₂ and He. At present, due to the computationally-intensive nature of numerically evaluating absorption strengths for molecular transitions and interactions for a given atmospheric species, specifying a grid of temperature and pressure values is necessary to obtain computable opacities for input into forward models and atmospheric retrieval algorithms. It is well known that extrapolating opacities above the temperature or pressure grid on which they were computed can be problematic (e.g. [77,78]), largely due to the dominance of transitions that originate in energy levels above the ground-state in higher temperature regimes. In this work we demonstrate that including self-broadening effects into the calculation of the absorption cross-section of water markedly affects the simulated transit depth of small-planet exoplanet atmospheres. Moreover, we are the first in the exoplanet field to illustrate that these generated differences are detectable with the near-future space missions JWST & Ariel, by explicitly simulating spectra with error bars for these instruments. We prove that self-broadening is necessary to account for in these calculations. As a community, our long-term goal should be to develop

cross-section functions, explicitly derived for pressure-temperature grids and as a function of molecular abundances, with all intra- and inter-molecular effects, like self-broadening, included.

5. Conclusion

In summary, it is evident that accounting for previously thought-to-be negligible absorption contributions, such as the self-broadening exhibited by H₂O in our opacity functions, will alter simulated transit depths by as much as 60 ppm. These differences sit above the noise level for a reasonable number of transit observations with the near-future space telescopes JWST and Ariel for the two small planets considered: GJ 1132 b and GJ 1214 b. Our quantification of the transit depth differences found by producing and utilising cross-sections which include the absorption contribution induced by H₂O self-broadening motivate further progress in not only refining such broadening parameters, but also developing opacities for a variety of molecular species expected to be found in the atmospheres of small planets, which may also be dominated by more than one heavy molecule. This is not an easy undertaking, due to the vast amount of work done and ongoing in the field of line shapes (including half widths $\gamma_{\text{H}_2\text{O}}$ and temperature dependence $n_{\text{H}_2\text{O}}$; see, for example, [41,42,44,55], and references therein). It is paramount that the field of exoplanet spectroscopy moves towards the use of more adaptive cross-sections, built as functions not only of temperature and pressure but also of molecular abundance, as we have illustrated specifically for the case of H₂O.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

CRediT authorship contribution statement

Lara O. Anisman: Methodology, Investigation, Software, Data curation, Formal analysis, Writing – original draft, Writing – review & editing, Visualization. **Katy L. Chubb:** Methodology, Software, Formal analysis, Writing – original draft, Writing – review & editing. **Quentin Changeat:** Conceptualization, Investigation, Supervision. **Billy Edwards:** Data curation, Software. **Sergei N. Yurchenko:** Conceptualization, Investigation, Supervision. **Jonathan Tennyson:** Conceptualization, Investigation, Writing – original draft, Writing – review & editing, Supervision, Funding acquisition. **Giovanna Tinetti:** Conceptualization, Investigation, Writing – original draft, Writing – review & editing, Supervision, Funding acquisition.

Acknowledgment

Funding: We acknowledge funding from the ERC through Consolidator grant ExoAI (GA 758892) and Advanced Grant Exo-MoHD (GA 883830), as well as from STFC grants ST/P000282/1, ST/P002153/1, ST/S002634/1 and ST/T001836/1. BE is a Laureate of the Paris Region fellowship programme which is supported by the Ile-de-France Region and has received funding under the Horizon 2020 innovation framework programme and the Marie Skłodowska-Curie grant agreement no. 945298.

References

- [1] de Wit J, Wakeford HR, Lewis NK, et al. Atmospheric reconnaissance of the habitable-zone earth-sized planets orbiting TRAPPIST-1. *Nat Astron* 2018;2(3):214–19. doi:10.1038/s41550-017-0374-z.
- [2] Tsiaras A, Rocchetto M, Waldmann IP, et al. Detection of an atmosphere around the super-Earth 55 Cancri e. *ApJ* 2016;820(2). arXiv:1511.08901. doi:10.3847/0004-637X/820/2/99. 99.

- [3] Demory B-O, Gillon M, de Wit J, Madhusudhan N, et al. A map of the large day–night temperature gradient of a super-earth exoplanet. *Nature* 2016;532(7598):207–9. doi:10.1038/nature17169.
- [4] Libby-Roberts JE, Berta-Thompson ZK, Diamond-Lowe H, et al. The featureless HST/WFC3 transmission spectrum of the rocky exoplanet GJ 1132b: no evidence for a cloud-free primordial atmosphere and constraints on starspot contamination. arXiv e-prints 2021. arXiv:2105.10487.
- [5] Mugnai LV, Modirrousta-Galian D, Edwards B, Changeat Q, et al. ARES V: no evidence for molecular absorption in the HST WFC3 spectrum of GJ 1132 b. arXiv e-prints 2021. arXiv:2104.01873.
- [6] Kreidberg L, Bean JL, Désert J-M, Benneke B, Deming D, Stevenson KB, et al. Clouds in the atmosphere of the super-Earth exoplanet GJ1214b. *Nature* 2014;505:69–72. arXiv:1401.0022. doi:10.1038/nature12888.
- [7] Tsiaras A, Waldmann IP, Tinetti G, Tennyson J, Yurchenko SN. Water vapour in the atmosphere of the habitable-zone eight-earth-mass planet k2-18 b. *Nat Astron* 2019. doi:10.1038/s41550-019-0878-9.
- [8] Edwards B, Changeat Q, Mori M, Anisman LO, et al. Hubble WFC3 spectroscopy of the habitable-zone super-earth LHS 1140 b. *Astron J (N Y)* 2020;161(1):44. doi:10.3847/1538-3881/abc6a5.
- [9] Owen JE, Wu Y. The evaporation valley in the kepler planets. *Apj* 2017;847. arXiv:1705.10810. doi:10.3847/1538-4357/aa890a. 29.
- [10] Fulton BJ, Petigura EA. The California kepler survey VII. precise planet radii leveraging gaia DR2 reveal the stellar mass dependence of the planet radius gap. ArXiv e-prints 2018. arXiv:1805.01453.
- [11] Van Eylen V, Agentoft C, Lundkvist MS, Kjeldsen H, Owen JE, Fulton BJ, et al. An asteroseismic view of the radius valley: stripped cores, not born rocky. *Mon Not R Astron Soc* 2018;479(4):4786–95. doi:10.1093/mnras/sty1783.
- [12] Valencia D, Ikoma M, Guillot T, Nettelmann N. Composition and fate of short-period super-earths. *Astron Astrophys* 2010;516:A20. doi:10.1051/0004-6361/200912839.
- [13] Lecante J, Forget F, Lammer H. On the (anticipated) diversity of terrestrial planet atmospheres. *Exp Astron* 2015;40:449–67. doi:10.1007/s10686-014-9403-4.
- [14] Dressing CD, Charbonneau D, Dumusque X, et al. The mass of kepler-93b and the composition of terrestrial planets. *Apj* 2015;800. arXiv:1412.8687. doi:10.1088/0004-637X/800/2/135. 135.
- [15] Valencia D, Guillot T, Parmentier V, Freedman RS. Bulk composition of GJ 1214b and other sub-neptune exoplanets. *Astrophys J* 2013;775(1):10. doi:10.1088/0004-637x/775/1/10.
- [16] Zeng L, Jacobsen SB, Sasselov DD, et al. Growth model interpretation of planet size distribution. *Proceedings of the National Academy of Sciences* 2019;116(20):9723–8. doi:10.1073/pnas.1812905116.
- [17] Tennyson J, Yurchenko SN, Al-Refaie AF, et al. The exomol database: molecular line lists for exoplanet and other hot atmospheres. *J Mol Spectrosc* 2016;327:73–94. doi:10.1016/j.jms.2016.05.002. *New Visions of Spectroscopic Databases, Volume II*.
- [18] Gordon I, Rothman LS, Wilzewski JS, Kochanov RV, Hill C, Tan Y, Wcislo P. HITRAN2016 : new and improved data and tools towards studies of planetary atmospheres. In: *AAS/Division for planetary sciences meeting abstracts #48*. In: *AAS/Division for Planetary Sciences Meeting Abstracts, vol. 48; 2016*. p. 421.13.
- [19] Bernath PF. MoLLIST: molecular line lists, intensities and spectra. *J Quant Spectrosc Radiat Transf* 2020;240:106687. doi:10.1016/j.jqsrt.2019.106687.
- [20] Rothman LS, Gordon IE, Barber RJ, et al. HITEMP, The high-temperature molecular spectroscopic database. *J Quant Spectrosc Radiat Transf* 2010;111:2139–50. doi:10.1016/j.jqsrt.2010.05.001.
- [21] Chubb KL, Rocchetto M, Al-Refaie AF, Waldmann I, Min M, Barstow J, et al. The exomolOP database: cross-sections and k-tables for molecules of interest in high-temperature exoplanet atmospheres. *A&A* 2020;646:A21.
- [22] Kitzmann D, Heng K, Oreshenko M, Grimm SL, Apai D, Bowler BP, et al. Helios-r2: a new bayesian, open-source retrieval model for brown dwarfs and exoplanet atmospheres. *Apj* 2020;890(2):174. doi:10.3847/1538-4357/ab6d71.
- [23] Phillips MW, Tremblin P, Baraffe I, Chabrier G, Allard NF, Spiegelman F, Goyal JM, Drummond B, Hebrard E. A new set of atmosphere and evolution models for cool t-y brown dwarfs and giant exoplanets. *A&A* 2020;637:A38. doi:10.1051/0004-6361/201937381.
- [24] Grimm SL, Malik M, Kitzmann D, et al. HELIOS-K 2.0 and an open-source opacity database for exoplanetary atmospheres. *The Astrophysical Journal Supplement* 2021;253:30. doi:10.3847/1538-4365/abd773.
- [25] Ducrot E, Gillon M, Delrez L, et al. Trappist-1: global results of the spitzer exploration science program red worlds. *Astronomy & Astrophysics* 2020;640:A112. doi:10.1051/0004-6361/201937392.
- [26] Crossfield IJM, Dragomir D, Cowan NB, Daylan T, Wong I, Kataria T, et al. Phase curves of hot neptune LTT 9779b suggest a high-metallicity atmosphere. *Astrophys J* 2020;903(1):L7. doi:10.3847/2041-8213/abc71.
- [27] Wunderlich F, Scheucher M, Godolt M, Grenfell JL, et al. Distinguishing between wet and dry atmospheres of TRAPPIST-1 e and f. *Astrophys J* 2020;901(2):126. doi:10.3847/1538-4357/aba59c.
- [28] Gharib-Nezhad E, Line MR. The influence of H₂O pressure broadening in high-metallicity exoplanet atmospheres. *Astrophys J* 2019;872(1):27. doi:10.3847/1538-4357/aabf7b.
- [29] Hedges C, Madhusudhan N. Effect of pressure broadening on molecular absorption cross sections in exoplanetary atmospheres. *Mon Not R Astron Soc* 2016;458:1427–49. doi:10.1093/mnras/stw278.
- [30] Rocchetto M. Characterisation of exoplanetary atmospheres and planetary systems. University College London, London, UK; 2017. Ph.D. thesis.
- [31] Barstow JK, Changeat Q, Garland R, Line MR, Rocchetto M, Waldmann IP. A comparison of exoplanet spectroscopic retrieval tools. *Mon Not R Astron Soc* 2020. doi:10.1093/mnras/staa548.
- [32] Gharib-Nezhad E, Iyer AR, Line MR, Freedman RS, Marley MS, Batalha NE. EXOPLINES: Molecular absorption cross-section database for brown dwarf and giant exoplanet atmospheres. *The Astrophysical Journal Supplement Series* 2021;254(2):34. doi:10.3847/1538-4365/abf504.
- [33] Anisman LO, Chubb KL, Eisey J, Al-Refaie A, Changeat Q, Yurchenko SN, et al. Cross-sections for heavy atmospheres: H₂O continuum. *J Quant Spectrosc Radiat Transfer* 2022;278:108013. doi:10.1016/j.jqsrt.2021.108013.
- [34] Skaf N, Bieger MF, Edwards B, Changeat Q, Morvan M, Kiefer F, et al. Ares. ii. characterizing the hot jupiters WASP-127 b, WASP-79 b, and WASP-62 b with the hubble space telescope. *Astron J (N Y)* 2020;160(3):109. doi:10.3847/1538-3881/ab94a3.
- [35] Pinhas A, Madhusudhan N, Gandhi S, MacDonald R. H₂O Abundances and cloud properties in ten hot giant exoplanets. *Mon Not R Astron Soc* 2019;482(2):1485–98. doi:10.1093/mnras/sty2544.
- [36] Sing DK, Fortney JJ, Nikolov N, et al. A continuum from clear to cloudy hot-jupiter exoplanets without primordial water depletion. *Nature* 2015;529(7584):59–62. doi:10.1038/nature16068.
- [37] Yurchenko SN, Al-Refaie AF, Tennyson J. EXOCROSS: A general program for generating spectra from molecular line lists. *A&A* 2018;614:A131. doi:10.1051/0004-6361/201732531.
- [38] Barton EJ, Hill C, Yurchenko SN, Tennyson J, Dudaryonok A, Lavrentieva NN. Pressure dependent water absorption cross sections for exoplanets and hot other atmospheres. *J Quant Spectrosc Radiat Transf* 2017;187:453–60. doi:10.1016/j.jqsrt.2016.10.024.
- [39] Barton EJ, Hill C, Czurylo M, Li H-Y, Hyslop A, Yurchenko SN, et al. The ExoMol diet of line-by-line pressure-broadening parameters. *J Quant Spectrosc Radiat Transf* 2017;203:490–5. doi:10.1016/j.jqsrt.2017.01.028.
- [40] Polyansky OL, Kyuberis AA, Zobov NF, Tennyson J, Yurchenko SN, Lodi L. Exomol molecular line lists XXX: a complete high-accuracy line list for water. *Mon Not R Astron Soc* 2018;480(2):2597–608.
- [41] Gamache RR, Hartmann J-M. An intercomparison of measured pressure-broadening and pressure-shifting parameters of water vapor. *Can J Chem* 2004;82:1013–27.
- [42] Gordon IE, Rothman LS, Hargreaves RJ, et al. The HITRAN2020 molecular spectroscopic database. *JQSRT* 2021;107949. doi:10.1016/j.jqsrt.2021.107949.
- [43] Gordon IE, Rothman LS, Gamache RR, et al. Current updates of the water-vapor line list in HITRAN: a new “diet” for air-broadened half-widths. *JQSRT* 2007;108(3):389–402. doi:10.1016/j.jqsrt.2007.06.009.
- [44] Toth RA. Measurements and analysis (using empirical functions for widths) of air- and self-broadening parameters of H₂O. *JQSRT* 2005;94(1):1–50. doi:10.1016/j.jqsrt.2004.08.041.
- [45] Loos J, Birk M, Wagner G. Measurement of positions, intensities and self-broadening line shape parameters of H₂O lines in the spectral ranges 1850–2280 cm⁻¹ and 2390–4000 cm⁻¹. *JQSRT* 2017;203:119–32. doi:10.1016/j.jqsrt.2017.02.013.
- [46] Tennyson J, Yurchenko SN, Al-Refaie AF, Clark VHJ, Chubb KL, et al. The 2020 release of the exomol database: molecular line lists for exoplanet and other hot atmospheres. *J Quant Spectrosc Radiat Transfer* 2020;255:107228. doi:10.1016/j.jqsrt.2020.107228.
- [47] Grossmann BE, Browell EV. Spectroscopy of water vapor in the 720-nm wavelength region: line strengths, self-induced pressure broadenings and shifts, and temperature dependence of linewidths and shifts. *J Mol Spectrosc* 1989;136(2):264–94. doi:10.1016/0022-2852(89)90336-6.
- [48] Markov VN. Temperature dependence of self-induced pressure broadening and shift of the 643-550 line of the water molecule. *J Mol Spectrosc* 1994;164(1):233–8. doi:10.1006/jmsp.1994.1069.
- [49] Bauer A, Godon M, Kheddar M, Hartmann JH, Bonamy J, Robert D. Temperature and perturber dependences of water-vapor 380 GHz-line broadening. *J Quant Spectrosc Radiat Transfer* 1987;37(6):531–9. doi:10.1016/0022-4073(87)90056-2.
- [50] Bauer A, Godon M, Kheddar M, Hartmann JM. Temperature and perturber dependences of water vapor line-broadening. experiments at 183 GHz; calculations below 1000 GHz. *JQSRT* 1989;41(1):49–54. doi:10.1016/0022-4073(89)90020-4.
- [51] Adler-Golden S, Lee J, Goldstein N. Diode laser measurements of temperature-dependent line parameters for water vapor near 820 nm. *JQSRT* 1992;48(5):527–35. doi:10.1016/0022-4073(92)90118-N. *Special Issue Conference on Molecular Spectroscopic Databases*.
- [52] Podobedov VB, Plusquellic DF, Fraser GT. The laser study of self-pressure and temperature broadening and shifts of water vapor lines for pressures up to 1.4kpa. *JQSRT* 2004;87(3):377–85. doi:10.1016/j.jqsrt.2004.03.001.
- [53] Gamache RR, Vispoel B. On the temperature dependence of half-widths and line shifts for molecular transitions in the microwave and infrared regions. *JQSRT* 2018;217:440–52. doi:10.1016/j.jqsrt.2018.05.019.
- [54] Wcislo P, Gordon IE, Tran H, et al. The implementation of non-voigt line profiles in the HITRAN database: H₂ case study. *J Quant Spectrosc Radiat Transf* 2016;177:75–91. doi:10.1016/j.jqsrt.2016.01.024. *XVIIIth Symposium on High Resolution Molecular Spectroscopy (HighRes-2015), Tomsk, Russia*.
- [55] Tan Y, Kochanov RV, Rothman LS, Gordon IE. Introduction of water-vapor broadening parameters and their temperature-dependent exponents into the HITRAN database: part I-CO₂, N₂O, CO, CH₄, O₂, NH₃, and H₂S. *Journal of Geophysical Research: Atmospheres* 2019;124(21):11580–94. doi:10.1029/2019JD030929.

- [56] Hartmann J-M, Tran H, Armante R, Boulet C, Campargue A, Forget F, Gianfrani L, Gordon I, Guerlet S, Gustafsson M, Hodges JT, Kassi S, Lisak D, Thibault F, Toon GC. Recent advances in collisional effects on spectra of molecular gases and their practical consequences. *J Quant Spectrosc Radiat Transf* 2018;213:178–227. doi:10.1016/j.jqsrt.2018.03.016.
- [57] Stolarczyk N, Thibault F, Cybulski H, Józwiak H, Kowzan G, Vispoel B, et al. Evaluation of different parameterizations of temperature dependences of the line-shape parameters based on ab initio calculations: case study for the HITRAN database. *J Quant Spectrosc Radiat Transf* 2020;240:106676. doi:10.1016/j.jqsrt.2019.106676.
- [58] Waldmann IP, Rocchetto M, Tinetti G, Barton EJ, Yurchenko SN, Tennyson J. Tau-REX II: retrieval of emission spectra. *ApJ* 2015;813. arXiv:1508.07591. doi:10.1088/0004-637X/813/1/13. 13.
- [59] Waldmann IP, Tinetti G, Rocchetto M, Barton EJ, Yurchenko SN, Tennyson J. Tau-REX I: A Next generation retrieval code for exoplanetary atmospheres. *ApJ* 2015;802. arXiv:1409.2312. doi:10.1088/0004-637X/802/2/107. 107.
- [60] Al-Refaei AF, Changeat Q, Waldmann IP, Tinetti G. TauREX III: a fast, dynamic and extendable framework for retrievals. arXiv e-prints 2019. arXiv:1912.07759.
- [61] Bonfils X, Almenara J-M, Cloutier R, Wünsche A, et al. Radial velocity follow-up of GJ1132 with HARPS. *Astronomy & Astrophysics* 2018;618:A142. doi:10.1051/0004-6361/201731884.
- [62] Harpsøe KBW, Hardis S, Hinse TC, Jørgensen UG, et al. The transiting system GJ1214: high-precision defocused transit observations and a search for evidence of transit timing variation. *A&A* 2013;549. arXiv:1207.3064. doi:10.1051/0004-6361/201219996. A10.
- [63] Abel M, Frommhold L, Li X, Hunt KLC. Collision-induced absorption by h2 pairs: from hundreds to thousands of kelvin. *The Journal of Physical Chemistry A* 2011;115(25):6805–12.
- [64] Fletcher LN, Gustafsson M, Orton GS. Hydrogen dimers in giant-planet infrared spectra. *The Astrophysical Journal Supplement Series* 2018;235(1):24.
- [65] Abel M, Frommhold L, Li X, Hunt KLC. Infrared absorption by collisional H₂-He complexes at temperatures up to 9000 K and frequencies from 0 to 20 000 cm⁻¹. *J Chem Phys* 2012;136(4):044319.
- [66] Anisman LO, Edwards B, Changeat Q, Venot O, Al-Refaei AF, Tsiaras A, et al. Wasp-117 b: an eccentric hot saturn as a future complex chemistry laboratory. *Astron J (N Y)* 2020;160(5):233. doi:10.3847/1538-3881/abb9b0.
- [67] Changeat Q, Edwards B, Waldmann IP, Tinetti G. Toward a more complex description of chemical profiles in exoplanet retrievals: a two-layer parameterization. *Astrophys J* 2019;886(1):39. doi:10.3847/1538-4357/ab4a14.
- [68] Rocchetto M, Waldmann IP, Venot O, Lagage P-O, Tinetti G. Exploring biases of atmospheric retrievals in simulated JWST transmission spectra of hot jupiters. *ApJ* 2016;833. arXiv:1610.02848. doi:10.3847/1538-4357/833/1/120. 120.
- [69] Tinetti G, Drossart P, Eccleston P, Hartogh P, Heske A, Leconte J, et al. A chemical survey of exoplanets with ARIEL. *Exp Astron* 2018;46(1):135–209. doi:10.1007/s10686-018-9598-x.
- [70] Tinetti G, Eccleston P, Haswell C, Lagage P-O, Leconte J, Lüftinger T, et al. Ariel: enabling planetary science across light-years. arXiv e-prints 2021. arXiv:2104.04824.
- [71] Edwards B, Mugnai L, Tinetti G, Pascale E, Sarkar S. An updated study of potential targets for ariel. *Astron J (N Y)* 2019;157(6):242. doi:10.3847/1538-3881/ab1cb9.
- [72] Gardner JP, Mather JC, Clampin M, et al. The james webb space telescope. *Space Sci Rev* 2006;123(4):485–606. arXiv:astro-ph/0606175. doi:10.1007/s11214-006-8315-7.
- [73] Cowan NB, Greene T, Angerhausen D, et al. Characterizing transiting planet atmospheres through 2025. *Publ Astron Soc Pac* 2015;127:311. arXiv:1502.00004. doi:10.1086/680855.
- [74] Mugnai LV, Pascale E, Edwards B, Papageorgiou A, Sarkar S. Arielrad: the ariel radiometric model. arXiv e-prints 2020. arXiv:2009.07824.
- [75] Edwards B, Stotesbury I. Terminus: A Versatile simulator for space-based telescopes. *AJ* 2021;161(6). arXiv:2101.10317. doi:10.3847/1538-3881/abdf4d. 266.
- [76] Pontoppidan KM, Pickering TE, Laidler VG, Gilbert K, et al. Pandeia: a multi-mission exposure time calculator for JWST and WFIRST. In: Peck AB, Seaman RL, Benn CR, editors. *Observatory operations: strategies, processes, and systems VI*. Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, vol. 9910; 2016. p. 991016. doi:10.1117/12.2231768.
- [77] Tinetti G, Beaulieu JP, Henning T, Meyer M, Micela G, Ribas I, et al. Echo. *Exp Astron* 2012;34(2):311–53. doi:10.1007/s10686-012-9303-4.
- [78] Yurchenko SN, Tennyson J, Bailey J, Hollis MDJ, Tinetti G. Spectrum of hot methane in astronomical objects using a comprehensive computed line list. *Proceedings of the National Academy of Sciences* 2014;111(26):9379–83. doi:10.1073/pnas.1324219111.
- [79] Hill C, Gordon IE, Kochanov RV, Barrett L, Rothman LS. HITRANonline: A modern online interface and the flexible representation of spectroscopic data in the HITRAN database. *J Quant Spectrosc Radiat Transf* 2016;177:4–14. doi:10.1016/j.jqsrt.2015.12.012.
- [80] Varanasi P, Chudamani S. The temperature dependence of lineshifts, linewidths and line intensities of methane at low temperatures. *J Quant Spectrosc Radiat Transf* 1990;43:1–11. doi:10.1016/0022-4073(90)90060-J.
- [81] Pine AS. Self-, N₂, O₂, H₂, Ar, and He broadening in the ν₃ band Q branch of CH₄. *J Chem Phys* 1992;97:773–85. doi:10.1063/1.463943.
- [82] Fox K, Jennings DE, Stern EA, Hunnard R. Measurements of argon-, helium-, hydrogen-, and nitrogen-broadened widths of methane lines near 9000 cm⁻¹. *J Quant Spectrosc Radiat Transf* 1998;39:473–6. doi:10.1016/0022-4073(88)90092-1.
- [83] Varanasi P, Tejwani GDT. Experimental and theoretical studies on collision-broadened lines in the ν₄-fundamental of methane. *J Quant Spectrosc Radiat Transf* 1972;12:849–55. doi:10.1016/0022-4073(72)90073-8.
- [84] Varanasi P, Chudamani S. Measurements of collision-broadened line widths in the ν₄-fundamental band of ¹²CH₄ at low temperatures. *J Quant Spectrosc Radiat Transf* 1989;41:335–43. doi:10.1016/0022-4073(89)90062-9.
- [85] Grigoriev IM, Filippov NN, Tonkov MV, Gabard T, Doucen RL. Estimation of line parameters under line mixing effects: the ν₃ band of CH₄ in helium. *J Quant Spectrosc Radiat Transf* 2001;69:189–204. doi:10.1016/S0022-4073(00)00076-5.
- [86] Gabard T, Grigoriev IM, Grigorovich NM, Tonkov MV. Helium and argon line broadening in the ν₂ band of CH₄. *J Mol Spectrosc* 2004;225:123–31. doi:10.1016/j.jms.2004.02.023.
- [87] Manne J, Bui TQ, Webster CR. Determination of foreign broadening coefficients for methane lines targeted by the tunable laser spectrometer (TLS) on the mars curiosity rover. *J Quant Spectrosc Radiat Transf* 2017;191:59–66. doi:10.1016/j.jqsrt.2017.01.035.
- [88] Gharib-Nezhad E, Heays AN, Bechtel HA, Lyons JR. H₂-induced pressure broadening and pressure shift in the P-branch of the ν₃ band of CH₄ from 300 to 655 K. *J Quant Spectrosc Radiat Transf* 2019;239:106649. doi:10.1016/j.jqsrt.2019.106649.
- [89] Faure A, Wiesenfeld L, Tennyson J, Drouin BJ. Pressure broadening of water and carbon monoxide transitions by molecular hydrogen at high temperatures. *J Quant Spectrosc Radiat Transf* 2013;116:79–86. doi:10.1016/j.jqsrt.2012.09.015.
- [90] Li G, Gordon IE, Rothman LS, Tan Y, Hu S-M, Kassi S, Campargue A, Medvedev ES. Rovibrational line lists for nine isotopologues of the CO molecule in the x¹Σ⁺ ground electronic state. *Astrophys J Suppl* 2015;216:15. doi:10.1088/0067-0049/216/1/15.
- [91] Mulvihill CR, Alturaiifi SA, Petersen EL. High-temperature He- and O₂-broadening of the R(12) line in the 1-0 band of carbon monoxide. *J Quant Spectrosc Radiat Transf* 2018;217:432–9. doi:10.1016/j.jqsrt.2018.06.015.
- [92] Mantz AW, Devi VM, Benner DC, Smith M, Predoi-Cross A, Dulick M. A multi-spectrum analysis of widths and shifts in the 2010–2260 cm⁻¹ region of 12c16o broadened by helium at temperatures between 80 and 297 K. *J Mol Struct* 2005;742(1–3):99–110.
- [93] Predoi-Cross A, Esteki K, Rozario H, Naseri H, Latif S, Thibault F, Devi VM, Smith M, Mantz AW. Theoretical and revisited experimentally retrieved he-broadened line parameters of carbon monoxide in the fundamental band. *J Quant Spectrosc Radiat Transf* 2016;184:322–40.
- [94] Sinclair PM, Duggan P, Berman R, Drummond JR, May AD. Line broadening in the fundamental band of CO in CO-he and CO-ar mixtures. *J Mol Spectrosc* 1998;191(2):258–64.
- [95] Voronin BA, Mishina TP, Lavrentyeva NN, Chesnokova TY, Zuev VE, Barber MJ, Tennyson J. Estimate of the J/J' dependence of water vapor line broadening parameters. *J Quant Spectrosc Radiat Transf* 2010;111:2308–14. doi:10.1016/j.jqsrt.2010.05.015.
- [96] Voronin BA, Lavrentyeva NN, Lugovskoy AA, Bykov AD, Starikov VI, Tennyson J. Self- and air-broadening coefficients of HD¹⁶O spectral lines. *Atmos Oceanic Opt* 2012;25(1):27–34. doi:10.1134/S1024856012010150.
- [97] Solodov AM, Starikov VI. Broadening and shift of lines of the ν₂ + ν₃ band of water vapor induced by helium pressure. *Opt Spectrosc* 2008;105(1):14–20. doi:10.1134/S0030400X08070035.
- [98] Solodov AM, Starikov VI. Helium-induced halfwidths and line shifts of water vapor transitions of the ν₁ + ν₂ and ν₂ + ν₃ bands. *Mol Phys* 2009;107(1):43–51. doi:10.1080/00268970802698655.
- [99] Petrova TM, Solodov AM, Solodov AA, Starikov VI. Vibrational dependence of an intermolecular potential for H₂O-he system. *J Quant Spectrosc Radiat Transf* 2013;129:241–53. doi:10.1016/j.jqsrt.2013.06.021.
- [100] Petrova TM, Solodov AM, Starikov VI, Solodov AA. Measurements and calculations of He-broadening and -shifting parameters of the transitions of the ν₁ + ν₂ + ν₃ bands. *Mol Phys* 2012;110:1493–503. doi:10.1080/00268976.2012.663939.
- [101] Petrova TM, Solodov AM, Solodov AA, Starikov VI. Broadening parameters of the H₂O-He collisional system for astrophysical applications. *J Mol Spectrosc* 2016;321:5–8. doi:10.1016/j.jms.2016.01.009.
- [102] Gamache RR, Vispoel B, Renaud CL, Cleghorn K, Hartmann L. Vibrational dependence, temperature dependence, and prediction of line shape parameters for the h₂O-H₂ collision system. *Icarus* 2019;326:186–96. doi:10.1016/j.icarus.2019.02.011.
- [103] Wilzewski JS, Gordon IE, Kochanov RV, Hill C, Rothman LS. H₂, He, and CO₂ line-broadening coefficients, pressure shifts and temperature-dependence exponents for the HITRAN database. part 1: SO₂, NH₃, HF, HCl, OCS and c₂h₂. *J Quant Spectrosc Radiat Transf* 2016;168:193–206. doi:10.1016/j.jqsrt.2015.09.003.